

Platinum(II) Complexes Bearing 1,1'-bis(diphenylphosphino)ferrocene as Building Blocks for Functionalized Redox Active Porphyrins

*Luigi Monsù Scolaro,^{*a} Maria Rosaria Plutino,^b Andrea Romeo,^a Raffaello Romeo,^a*

Giampaolo Ricciardi,^c Sandra Belviso,^c Alberto Albinati,^d

Electronic Supporting Information

Fig. ESI1. $^{31}\text{P}\{^1\text{H}\}$ NMR spectra recorded in CD_2Cl_2 at 298 K relative to the complexes $[\text{PtMeCl}(\text{dppf})]$ (A), $[\text{PtMe}(\text{Me}_2\text{SO})(\text{dppf})]\text{CF}_3\text{SO}_3$ (B), and $[\text{PtMe}(\text{dppf})]_4\text{TpyP}(\text{CF}_3\text{SO}_3)_4$ (C).

Fig. ESI2. Absorption spectra of samples obtained by evaporating dichloromethane solutions of complexes **3** and **5** on glass cover slides.

Fig. ESI3. Fluorescence spectra of samples obtained by evaporating dichloromethane solutions of complexes **3** and **5** on glass cover slides.




Fig. ESI1




Fig. ESI2




Fig. ESI3