

Electronic Supplementary Information File

**Multiwalled carbon nanotubes-agarose gel micro-solid phase extraction for the
determination of triazine herbicides in water samples**

Mohd Marsin Sanagi^{a,b,*}, Mohamad Nazirul Muzakkir Mat Jais^a, Sazlinda Kamaruzaman^{a,c}, Wan

Aini Wan Ibrahim^a, Umair Baig^{a,b}

^aDepartment of Chemistry, Faculty of Science, Universiti Teknologi Malaysia,

81310 UTM Johor Bahru, Johor, Malaysia

^bIbnu Sina Institute for Fundamental Science Studies, Nanotechnology Research Alliance,

Universiti Teknologi Malaysia, 81310 UTM Johor Bahru, Johor, Malaysia

^cDepartment of Chemistry, Faculty of Science, Universiti Putra Malaysia,

43400 UPM Serdang Selangor, Malaysia

**Corresponding author, E-mail: marsin@kimia.fs.utm.my (M. M. Sanagi)*

Tel: +607-5534517 Fax: +607-5566162




Fig. S1. The chemical structures of (a) ATR and (b) SEC.




Fig. S2. Chromatogram of non-spiked tap water (a), spiked tap water at 5 ppb (b), and spiked tap water at 100 ppb (c) after MWCNTs-AG- μ SPE under the optimum extraction conditions. GC conditions as in Fig. 6.