

Electronic Supplementary Material (ESI) for CrystEngComm.
This journal is © The Royal Society of Chemistry 2015

Electronic Supplementary Information

The first photochromic bimetallic assemblies based on Mn(III) and Mn(II) Schiff-base (salpn, dapsc) complexes and pentacyanonitrosylferrate

Vyacheslav A. Kopotkov,^{a*} Valentina D. Sasnovskaya,^a Denis V. Korchagin,^a Roman B. Morgunov,^a Sergey M. Aldoshin,^a Sergey V. Simonov,^b Leokadiya V. Zorina,^{b*} Dominik Schaniel,^{c,d} Theo Woike,^{e*} and Eduard B. Yagubskii^{a*}

^a Institute of Problems of Chemical Physics, Chernogolovka, Moscow District, 1 Academician Semenov av., 142432 Russian Federation. E-mail: yagubski@icp.ac.ru, slavaoven@mail.ru; Fax: +7 496-5225636; Tel: +7 495-9935707

^b Institute of Solid State Physics, Chernogolovka, Moscow District, 2 Academician Ossipyan str., 142432 Russian Federation. E-mail: zorina@issp.ac.ru; Fax: +7 496-5228160; Tel: +7 496-5221982

^c Université de Lorraine, CRM2, UMR 7036, Vandoeuvre-les-Nancy, F-54506, France.
E-mail: dominik.schaniel@univ-lorraine.fr; Fax: +33 (0) 383 68 4300; Tel: +33 (0) 383 68 4863

^d CNRS, CRM², UMR 7036, Vandoeuvre-les-Nancy, F-54506, France,. E-mail: dominik.schaniel@univ-lorraine.fr;

^e Technische Universität Dresden, Institut für Strukturphysik, Zellescher Weg 16, 01069 Dresden, Germany.
E-mail: theo.woike@tu-dresden.de Fax: +49 351 463-37048; Tel: +49 351 463-34670




Fig. S1 TG-DSC curves and mass spectra for complex 1.




Fig. S2 TG-DSC curves and mass spectra for freshly prepared complex 3.

Crystal structures




Fig. S3 Structure of $\{[\text{Mn}(\text{salpn})(\text{CH}_3\text{OH})]_4[\text{Fe}(\text{CN})_5\text{NO}]\}(\text{ClO}_4)_2 \cdot 4\text{H}_2\text{O}$ (**4**) (space group P2(1)/n, $a=15.372(1)\text{\AA}$, $b=17.519(1)\text{\AA}$, $c=15.692(1)\text{\AA}$, $\beta=90.05(1)^\circ$). H atoms are omitted for clarity.

As in the case of compound **2**, the nitrosyl group is disordered between two sites with 1/2 occupancies due to space group symmetry.




Fig. S4 The environment of NO ligand in complex $\{[\text{Mn}(\text{salpn})(\text{CH}_3\text{OH})]_4[\text{Mn}(\text{CN})_5\text{NO}]\}[\text{C}(\text{CN})_3] \cdot 3\text{H}_2\text{O}$ (**2**) (the distances in \text{\AA}).