

Supporting Information

Utilizing alkoxyphenyl substituents for side-chain engineering of efficient benzo[1,2-*b*:4,5-*b'*]dithiophene-based small molecule organic solar cells

*Zhengkun Du^{a,c}, Weichao Chen^a, Meng Qiu^a, Yanhua Chen^{a,b}, Ning Wang^a, Ting Wang^a,
Mingliang Sun^{*b}, Donghong Yu^{*c,d}, and Renqiang Yang^{*a}*

^a CAS Key Laboratory of Bio-based Materials, Qingdao Institute of Bioenergy and Bioprocess Technology, Chinese Academy of Sciences, Qingdao 266101, China,

^b Institute of Materials Science and Engineering, Ocean University of China, Qingdao 266100, China,

^c Department of Chemistry and Bioscience, Aalborg University, DK-9220, Aalborg, Denmark

^d Sino-Danish Centre for Education and Research (SDC), Niels Jensens Vej 2, DK-8000, Aarhus, Denmark.




Figure S1. DSC thermograms of DCA3TBDTP.




Figure S2. UV-vis absorption spectrum of DCA3TBDTP in CHCl₃ solution (5×10^{-6} mol/L).




Figure S3. (a) ^1H NMR and (b) ^{13}C NMR spectra of DCA3TBDTP.