

Influence of selected inorganic counter-ions on the structure and antimicrobial properties of silver(I) complexes with imidazole-containing ligand


Urszula Kalinowska-Lis,^{a,*} Aleksandra Felczak,^b Lilianna Chęcińska,^c Magdalena Małecka,^c Katarzyna Lisowska,^b and Justyn Ochocki^a

Electronic Supplementary Information (ESI)
for New Journal of Chemistry


Fig. S2

Fig. S2 ^1H NMR(D_2O) spectra of the free ligand 4(5)- CH_2OHimH and silver(I) complexes **1-5**


^1H NMR(D_2O) spectrum of free ligand 4(5)- CH_2OHimH (peak at 4,71 ppm is from D_2O)




^1H NMR(D_2O) spectrum of $[\text{Ag}(4\text{-CH}_2\text{OHimH})_2]\text{NO}_3$ (**1**)




^1H NMR(D_2O) spectrum of $[\text{Ag}(\text{4-CH}_2\text{OHimH})_2]\text{ClO}_4$ (**2**)



^1H NMR(D_2O) spectrum of $[\text{Ag}(\text{4-CH}_2\text{OHimH})_2]\text{CF}_3\text{COO}$ (**3**)



^1H NMR(D_2O) spectrum of $[\text{Ag}(\text{4-CH}_2\text{OHimH})_2]\text{BF}_4$ (**4**)



^1H NMR(D_2O) spectrum of $[\text{Ag}(\text{4-CH}_2\text{OHimH})_2]\text{SO}_3\text{CH}_3$ (**5**)

