

Supplementary Information

Hyperbranched fluorene-alt-carbazole copolymers with spiro[3.3]heptane-2,6-dispirofluorene as the core and their application in white polymer light-emitting devices

Yuling Wu,^{ab} Jie Li,^{ab} Wenqing Liang,^{ab} Junli Yang,^{ab} Jing Sun,^{ab} Hua Wang,^{ab}
Xuguang Liu,^c Bingshe Xu,^{*ab} Wei Huang,^d

^a Key Laboratory of Interface Science and Engineering in Advanced Materials, Taiyuan University of Technology, Taiyuan, 030024, China.

^b Research Center of Advanced Materials Science and Technology, Taiyuan University of Technology, Taiyuan, 030024, China.

^c College of Chemistry and Chemical Engineering, Taiyuan University of Technology, Taiyuan, 030024, China.

^d Key Laboratory for Organic Electronics & Information Displays (KLOEID) and Institute of Advanced Materials, Nanjing University of Posts and Telecommunications (NUPT), Nanjing, 210046, China.

* Corresponding author. E-mail: Bingshe Xu: xubs@tyut.edu.cn




Figure S1. ^1H NMR spectrum of DBrCz.




Figure S2. ^{13}C NMR spectrum of DBrCz.




Figure S3. ^1H NMR spectra of the copolymers.




Figure S4. ^1H NMR spectrum of PFCzSDF10DBT7.





Figure S5. UV-vis absorption of DBT and PL spectra of **PFCz** in CHCl_3 solution (10^{-5} mol/L) (a), and PL spectra of **DBT** in CHCl_3 solution (10^{-5} mol/L) (b).