

**Supporting Information**

**For**

**Selective removal of nitroaromatic compounds from wastewater in an integrated  
zero valent iron (ZVI) reduction and ZVI/H<sub>2</sub>O<sub>2</sub> oxidation process**

Jianguo Liu<sup>1</sup>, Changjin Ou<sup>1</sup>, Weiqing Han, Faheem, Jinyou Shen\*, Huiping Bi,

Xiuyun Sun, Jiansheng Li, Lianjun Wang\*


*Jiangsu Key Laboratory for Chemical Pollution Control and Resources Reuse, School  
of Environmental and Biological Engineering, Nanjing University of Science and  
Technology, Nanjing, 210094, China*

Corresponding author: \*Jinyou Shen, Tel/Fax: +86 25 84303965, E-mail address:  
shenjinyou@mail.njust.edu.cn; \*Lianjun Wang, Tel/Fax: +86 25 84315941, E-mail  
address: wanglj@mail.njust.edu.cn.


<sup>1</sup>These authors contributed to the paper equally.

**Table S1** Characteristics of DNAN producing wastewater


Parameter	Value
pH	7.2
COD (mg/L)	34000±985
TOC (mg/L)	7151±329
DNAN (mg/L)	123.1±3.8
DNCB (mg/L)	249.3±6.9
DNP (mg/L)	110.4±4.2
Salinity (%)	3.3
EC <sub>50, 48 h (v/v)</sub> (%)	0.67



**Fig.S1** SEM images of fresh ZVI (a), ZVI shavings after reduction (b) and ZVI shavings after oxidation (c)



**Fig.S2** Evolution of UV-vis spectra before and after treatment



**Fig.S3** Evolution of HPLC spectra before and after treatment