

Supplementary Information

Sulfonated nanohydroxyapatite functionalized with 2-aminoethyl dihydrogen phosphate (HAP@AEPH₂-SO₃H) as a new recyclable and eco-friendly catalyst for rapid one-pot synthesis of 4,4'-(aryl methylene)bis(3-methyl-1*H*-pyrazol-5-ol)s

Monireh Zarghani, Batool Akhlaghinia*

*Department of Chemistry, Faculty of Sciences, Ferdowsi University of Mashhad, Mashhad 9177948974,
Iran.*

Experimental

General

The purity determinations of the products were accomplished by TLC on silica gel polygram STL G/UV 254 plates. The melting points of products were determined with an Electrothermal Type 9100 melting point apparatus. The FTIR spectra were recorded on pressed KBr pellets using a AVATAR 370 FT-IR spectrometer (Therma Nicolet spectrometer, USA) at room temperature in the range between 4000 and 400 cm⁻¹ with a resolution of 4 cm⁻¹, and each spectrum was the average of 32 scans. NMR spectra were recorded on a NMR Bruker Avance spectrometer at 400 and 300 MHz in DMSO-d₆ as solvent in the presence of tetramethylsilane as internal standard. Mass spectra were recorded with a CH7A Varianmat Bremem instrument at 70

eV electron impact ionization; in m/z (rel %). All of the products were known compounds and characterized by the FT-IR and comparison of their melting points with known compounds. The structure of selected products was further confirmed by ^1H NMR, ^{13}C NMR spectroscopy, mass spectrometry and elemental analysis.

4,4'-(Phenylmethylene)bis(3-methyl-1-phenyl-1*H*-pyrazol-5-ol)(3a)




Figure 1a: ¹H NMR (300 MHz, DMSO-*d*₆) of 4,4'-(Phenylmethylene)bis(3-methyl-1-phenyl-1*H*-pyrazol-5-ol)




Figure 1b: ^{13}C NMR (75 MHz, DMSO- d_6) of 4,4'-(Phenylmethylene)bis(3-methyl-1-phenyl-1*H*-pyrazol-5-ol)




Figure 1c: Mass spectrum of 4,4'-(Phenylmethylene)bis(3-methyl-1-phenyl-1*H*-pyrazol-5-ol).





Figure 1d: FT-IR (KBr) spectrum of 4,4'-(Phenylmethylene)bis(3-methyl-1-phenyl-1*H*-pyrazol-5-ol).

4,4'-(4-Hydroxyphenyl)methylene)bis(3-methyl-1-phenyl-1*H*-pyrazol-5-ol) (3b)





Figure 2a: ^1H NMR (300 MHz, DMSO- d_6) of 4,4'-(4-Hydroxyphenyl)methylene)bis(3-methyl-1-phenyl-1*H*-pyrazol-5-ol)




Figure 2b: ^{13}C NMR (75 MHz, DMSO- d_6) of 4,4'-(4-Hydroxyphenyl)methylene)bis(3-methyl-1-phenyl-1*H*-pyrazol-5-ol)




Figure 2c: Mass spectrum of 4,4'-(4-Hydroxyphenyl)methylene)bis(3-methyl-1-phenyl-1*H*-pyrazol-5-ol)




Figure 2d: FT-IR (KBr) spectrum of 4,4'-(4-Hydroxyphenyl)methylene)bis(3-methyl-1-phenyl-1*H*-pyrazol-5-ol).

4,4'-(*(3*-Hydroxyphenyl)methylene)bis(*3*-methyl-1-phenyl-1*H*-pyrazol-5-ol) (3c)




Figure 3a: ¹H NMR (300 MHz, DMSO-*d*₆) of 4,4'-(*(3*-Hydroxyphenyl)methylene)bis(*3*-methyl-1-phenyl-1*H*-pyrazol-5-ol)




Figure 3b: ^{13}C NMR (75 MHz, DMSO- d_6) of 4,4'-(3-Hydroxyphenyl)methylenebis(3-methyl-1-phenyl-1*H*-pyrazol-5-ol)




Figure 3c: FT-IR (KBr) spectrum of 4,4'-(3-Hydroxyphenyl)methylenebis(3-methyl-1-phenyl-1*H*-pyrazol-5-ol).

4,4'-(*(2*-Hydroxyphenyl)methylene)bis(3-methyl-1-phenyl-1*H*-pyrazol-5-ol) (3d)




Figure 4a: ¹H NMR (300 MHz, DMSO-*d*₆) of 4,4'-(*(2*-Hydroxyphenyl)methylene)bis(3-methyl-1-phenyl-1*H*-pyrazol-5-ol)




Figure 4b: Mass spectrum of 4,4'-(2-Hydroxyphenyl)methylene)bis(3-methyl-1-phenyl-1*H*-pyrazol-5-ol).




Figure 4c: FT-IR (KBr) spectrum of 4,4'-(2-Hydroxyphenyl)methylene)bis(3-methyl-1-phenyl-1*H*-pyrazol-5-ol).

4,4'-(*(4*-Methoxyphenyl)methylene)bis(*3*-methyl-1-phenyl-1*H*-pyrazol-5-ol)) (3e**)**




Figure 5a: ¹H NMR (300 MHz, DMSO-*d*₆) of 4,4'-(*(4*-Methoxyphenyl)methylene)bis(*3*-methyl-1-phenyl-1*H*-pyrazol-5-ol))




Figure 5b: ^{13}C NMR (75 MHz, DMSO- d_6) of 4,4'-(4-Methoxyphenyl)methylene)bis(3-methyl-1-phenyl-1*H*-pyrazol-5-ol))




Figure 5c: Mass spectrum of 4,4'-(4-Methoxyphenyl)methylene)bis(3-methyl-1-phenyl-1*H*-pyrazol-5-ol)).




Figure 5d: FT-IR (KBr) spectrum of 4,4'-(4-Methoxyphenyl)methylenebis(3-methyl-1-phenyl-1*H*-pyrazol-5-ol))

4,4'-(*(3*-Methoxyphenyl)methylene)bis(*3*-methyl-1-phenyl-1*H*-pyrazol-5-ol) (3f**)**




Figure 6a: ¹H NMR (300 MHz, DMSO-*d*₆) of 4,4'-(*(3*-Methoxyphenyl)methylene)bis(*3*-methyl-1-phenyl-1*H*-pyrazol-5-ol)




Figure 6b: Mass spectrum of 4,4'-(3-Methoxyphenyl)methylene)bis(3-methyl-1-phenyl-1*H*-pyrazol-5-ol).




Figure 6c: FT-IR (KBr) spectrum of 4,4'-(3-Methoxyphenyl)methylene)bis(3-methyl-1-phenyl-1*H*-pyrazol-5-ol).

4,4'-(*p*-Tolylmethylene)bis(3-methyl-1-phenyl-1*H*-pyrazol-5-ol)(3g)




Figure 7a: ¹H NMR (300 MHz, DMSO-*d*₆) of 4,4'-(*p*-Tolylmethylene)bis(3-methyl-1-phenyl-1*H*-pyrazol-5-ol)




Figure 7b: ^{13}C NMR (75 MHz, $\text{DMSO}-d_6$) of 4,4'-(*p*-Tolylmethylene)bis(3-methyl-1-phenyl-1*H*-pyrazol-5-ol)




Figure 7c: Mass spectrum of 4,4'-(*p*-Tolylmethylene)bis(3-methyl-1-phenyl-1*H*-pyrazol-5-ol).




Figure 7d: FT-IR (KBr) spectrum of 4,4'-(*p*-Tolylmethylene)bis(3-methyl-1-phenyl-1*H*-pyrazol-5-ol).

4,4'-(*m*-Tolylmethylene)bis(3-methyl-1-phenyl-1*H*-pyrazol-5-ol) (3h)




Figure 8a: ¹H NMR (300 MHz, DMSO-*d*₆) of 4,4'-(*m*-Tolylmethylene)bis(3-methyl-1-phenyl-1*H*-pyrazol-5-ol).




Figure 8b: ^{13}C NMR (75 MHz, DMSO- d_6) of 4,4'-(*m*-Tolylmethylene)bis(3-methyl-1-phenyl-1*H*-pyrazol-5-ol).




Figure 8c: Mass spectrum of 4,4'-(*m*-Tolylmethylene)bis(3-methyl-1-phenyl-1*H*-pyrazol-5-ol).




Figure 8d: FT-IR (KBr) spectrum of 4,4'-(*m*-Tolylmethylene)bis(3-methyl-1-phenyl-1*H*-pyrazol-5-ol).

4,4'-(5-Bromo-2-hydroxyphenyl)methylene)bis(3-methyl-1-phenyl-1*H*-pyrazol-5-ol) (3i**)**




Figure 9a: ¹H NMR (400 MHz, DMSO-*d*₆) of 4,4'-(5-Bromo-2-hydroxyphenyl)methylene)bis(3-methyl-1-phenyl-1*H*-pyrazol-5-ol).




Figure 9b: ^{13}C NMR (75 MHz, DMSO- d_6) of 4,4'-(5-Bromo-2-hydroxyphenyl)methylene)bis(3-methyl-1-phenyl-1*H*-pyrazol-5-ol).




Figure 9c: FT-IR (KBr) spectrum of 4,4'-(5-Bromo-2-hydroxyphenyl)methylene)bis(3-methyl-1-phenyl-1*H*-pyrazol-5-ol).

4,4'-(*(4*-Isopropylphenyl)methylene)bis(3-methyl-1-phenyl-1*H*-pyrazol-5-ol) (3j)




Figure 10a: ¹H NMR (300 MHz, DMSO-*d*₆) of 4,4'-(4-isopropylphenyl)methylenebis(3-methyl-1-phenyl-1*H*-pyrazol-5-ol)




Figure 10b: Mass spectrum of 4,4'-(4-isopropylphenyl)methylenebis(3-methyl-1-phenyl-1*H*-pyrazol-5-ol).




Figure 10c: FT-IR (KBr) spectrum of 4,4'-(4-isopropylphenyl)methylenebis(3-methyl-1-phenyl-1*H*-pyrazol-5-ol).

4,4'-(2-Chlorophenyl)methylene)bis(3-methyl-1-phenyl-1*H*-pyrazol-5-ol) (3k)




Figure 11a: ¹H NMR (300 MHz, DMSO-*d*₆) of 4,4'-(2-Chlorophenyl)methylene)bis(3-methyl-1-phenyl-1*H*-pyrazol-5-ol)




Figure 11b: ^{13}C NMR (75 MHz, $\text{DMSO}-d_6$) of 4,4'-(2-Chlorophenyl)methylenebis(3-methyl-1-phenyl-1*H*-pyrazol-5-ol).




Figure 11c: Mass spectrum of 4,4'-(2-Chlorophenyl)methylenebis(3-methyl-1-phenyl-1*H*-pyrazol-5-ol).




Figure 11d: FT-IR (KBr) spectrum of 4,4'-(2-Chlorophenyl)methylene)bis(3-methyl-1-phenyl-1*H*-pyrazol-5-ol).

4,4'-(*(4*-Chlorophenyl)methylene)bis(3-methyl-1-phenyl-1*H*-pyrazol-5-ol) (3l**)**




Figure 12a: ¹H NMR (300 MHz, DMSO-*d*₆) of 4,4'-(4-Chlorophenyl)methylene)bis(3-methyl-1-phenyl-1*H*-pyrazol-5-ol).




Figure 12b: ^{13}C NMR (75 MHz, DMSO- d_6) of 4,4'-(4-Chlorophenyl)methylene)bis(3-methyl-1-phenyl-1*H*-pyrazol-5-ol).





Figure 12c: Mass spectrum of 4,4'-(4-chlorophenyl)methylenebis(3-methyl-1-phenyl-1*H*-pyrazol-5-ol).






Figure 12d: FT-IR (KBr) spectrum of 4,4'-(4-Chlorophenyl)methylenebis(3-methyl-1-phenyl-1*H*-pyrazol-5-ol).

4,4'-(*(2,4-Dichlorophenyl)methylene*)bis(3-methyl-1-phenyl-1*H*-pyrazol-5-ol) (3m)




Figure 13a: ¹H NMR (300 MHz, DMSO-*d*₆) of 4,4'-(*(2,4-Dichlorophenyl)methylene*)bis(3-methyl-1-phenyl-1*H*-pyrazol-5-ol).




Figure 13b: ^{13}C NMR (75 MHz, DMSO-*d*₆) of 4,4'-(2,4-Dichlorophenyl)methylene)bis(3-methyl-1-phenyl-1*H*-pyrazol-5-ol).




Figure 13c: FT-IR (KBr) spectrum of 4,4'-(2,4-Dichlorophenyl)methylene)bis(3-methyl-1-phenyl-1*H*-pyrazol-5-ol).

4,4'-(*(4*-Bromophenyl)methylene)bis(3-methyl-1-phenyl-1*H*-pyrazol-5-ol) (3n)




Figure 14a: ¹H NMR (400 MHz, DMSO-*d*₆) of 4,4'-(*(4*-Bromophenyl)methylene)bis(3-methyl-1-phenyl-1*H*-pyrazol-5-ol).




Figure 14b: FT-IR (KBr) spectrum of 4,4'-(4-Bromophenyl)methylene)bis(3-methyl-1-phenyl-1*H*-pyrazol-5-ol).

4,4'-(*(3*-Nitrophenyl)methylene)bis(3-methyl-1-phenyl-1*H*-pyrazol-5-ol) (3o**)**




Figure 15a: ¹H NMR (300 MHz, DMSO-*d*₆) of 4, 4'-(*(3*-Nitrophenyl)methylene)bis(3-methyl-1-phenyl-1*H*-pyrazol-5-ol)





Figure 15b: Mass spectrum of 4, 4'-(3-Nitrophenyl)methylene)bis(3-methyl-1-phenyl-1*H*-pyrazol-5-ol).






Figure 15c: FT-IR (KBr) spectrum of 4,4'-(3-Nitrophenyl)methylene)bis(3-methyl-1-phenyl-1*H*-pyrazol-5-ol).

4,4'-(2-Nitrophenyl)methylene)bis(3-methyl-1-phenyl-1*H*-pyrazol-5-ol) (3p)




Figure 16a: ¹H NMR (300 MHz, DMSO-*d*₆) of 4, 4'-(2-Nitrophenyl)methylene)bis(3-methyl-1-phenyl-1*H*-pyrazol-5-ol).




Figure 16b: ^{13}C NMR (75 MHz, $\text{DMSO}-d_6$) of 4, 4'-(2-Nitrophenyl)methylene)bis(3-methyl-1-phenyl- $1H$ -pyrazol-5-ol)




Figure 16c: Mass spectrum of 4,4'-(2-Nitrophenyl)methylenebis(3-methyl-1-phenyl-1*H*-pyrazol-5-ol).




Figure 16d: FT-IR (KBr) spectrum of 4,4'-(2-Nitrophenyl)methylenebis(3-methyl-1-phenyl-1*H*-pyrazol-5-ol).

4,4'-(Naphthalen-1-ylmethylene)bis(3-methyl-1-phenyl-1*H*-pyrazol-5-ol) (3q)




Figure 17a: ¹H NMR (400 MHz, DMSO-*d*₆) of 4,4'-(Naphthalen-1-ylmethylene)bis(3-methyl-1-phenyl-1*H*-pyrazol-5-ol)




Figure 17b: FT-IR (KBr) spectrum of 4,4'-(Naphthalen-1-ylmethylene)bis(3-methyl-1-phenyl-1*H*-pyrazol-5-ol).

4,4'-(Pyridin-2-ylmethylene)bis(3-methyl-1-phenyl-1*H*-pyrazol-5-ol) (3r)




Figure 18a: ¹H NMR (300 MHz, DMSO-*d*₆) of 4,4'-(Pyridin-2-ylmethylene)bis(3-methyl-1-phenyl-1*H*-pyrazol-5-ol)




Figure 18b: ^{13}C NMR (75 MHz, DMSO- d_6) of 4,4'-(Pyridin-2-ylmethylene)bis(3-methyl-1-phenyl-1*H*-pyrazol-5-ol)




Figure 18c: FT-IR (KBr) spectrum of 4,4'-(Pyridin-2-ylmethylene)bis(3-methyl-1-phenyl-1*H*-pyrazol-5-ol)

4,4'-(Pyridin-3-ylmethylene)bis(3-methyl-1-phenyl-1*H*-pyrazol-5-ol) (3s)




Figure 19a: ¹H NMR (300 MHz, DMSO-*d*₆) of 4,4'-(Pyridin-3-ylmethylene)bis(3-methyl-1-phenyl-1*H*-pyrazol-5-ol).




Figure 19b: ^{13}C NMR (75MHz, DMSO- d_6) of 4,4'-(Pyridin-3-ylmethylene)bis(3-methyl-1-phenyl-1*H*-pyrazol-5-ol).





Figure 19c: Mass spectrum of **4,4'-(Pyridin-3-ylmethylene)bis(3-methyl-1-phenyl-1*H*-pyrazol-5-ol)**






Figure 19d: FT-IR (KBr) spectrum of 4,4'-(Pyridin-3-ylmethylene)bis(3-methyl-1-phenyl-1*H*-pyrazol-5-ol).

(E)-4,4'-(3-Phenylprop-2-ene-1,1-diyl)bis(3-methyl-1-phenyl-1*H*-pyrazol-5-ol) (3t)




Figure 20a: ¹H NMR (300 MHz, DMSO-*d*₆) of (*E*)-4,4'-(3-Phenylprop-2-ene-1,1-diyl)bis(3-methyl-1-phenyl-1*H*-pyrazol-5-ol)




Figure 20b: ^{13}C NMR (75 MHz, $\text{DMSO}-d_6$) of of (*E*)-4,4'-(3-Phenylprop-2-ene-1,1-diyl)bis(3-methyl-1-phenyl-1*H*-pyrazol-5-ol)




Figure 20c: Mass spectrum of (*E*)-4,4'-(3-Phenylprop-2-ene-1,1-diyl)bis(3-methyl-1-phenyl-1*H*-pyrazol-5-ol)





Figure 20d: FT-IR (KBr) spectrum of (*E*)-4,4'-(3-Phenylprop-2-ene-1,1-diyl)bis(3-methyl-1-phenyl-1*H*-pyrazol-5-ol).

4,4'-(Phenylmethylene)bis(3-methyl-1*H*-pyrazol-5-ol) (3u)




Figure 21a: ¹H NMR (300 MHz, DMSO-*d*₆) of 4,4'-(Phenylmethylene)bis(3-methyl-1*H*-pyrazol-5-ol).





Figure 21b: Mass spectrum of 4,4'-(Phenylmethylene)bis(3-methyl-1*H*-pyrazol-5-ol).






Figure 21c: FT-IR (KBr) spectrum of 4,4'-(Phenylmethylene)bis(3-methyl-1*H*-pyrazol-5-ol).

4,4'-(4-Chlorophenyl)methylene)bis(3-methyl-1*H*-pyrazol-5-ol) (3v)




Figure 22a: ¹H NMR (300 MHz, DMSO-*d*₆) of 4,4'-(4-Chlorophenyl)methylene)bis(3-methyl-1*H*-pyrazol-5-ol).




Figure 22b: Mass spectrum of 4,4'-(4-Chlorophenyl)methylene)bis(3-methyl-1*H*-pyrazol-5-ol).




Figure 22c: FT-IR (KBr) spectrum of 4,4'-(4-Chlorophenyl)methylene)bis(3-methyl-1*H*-pyrazol-5-ol).

4,4'-(*p*-Tolylmethylene)bis(3-methyl-1*H*-pyrazol-5-ol) (3w)





Figure 23a: ^1H NMR (300 MHz, $\text{DMSO}-d_6$) of 4,4'-(*p*-Tolylmethylene)bis(3-methyl-1*H*-pyrazol-5-ol).




Figure 23b: ^{13}C NMR (75 MHz, $\text{DMSO}-d_6$) of 4,4'-(*p*-Tolylmethylene)bis(3-methyl-1*H*-pyrazol-5-ol)




Figure 23c: Mass spectrum of 4,4'-(*p*-Tolylmethylene)bis(3-methyl-1*H*-pyrazol-5-ol).




Figure 23 d: FT-IR (KBr) spectrum of 4,4'-(*p*-Tolylmethylene)bis(3-methyl-1*H*-pyrazol-5-ol).

(E)-4-(4-(Dimethylamino)benzylidene)-3-methyl-1-phenyl-1*H*-pyrazol-5(4*H*)-one




Figure 24a: ¹H NMR (300 MHz, DMSO-*d*₆) of (E)-4-(4-(Dimethylamino)benzylidene)-3-methyl-1-phenyl-1*H*-pyrazol-5(4*H*)-one.




Figure 24b: Mass spectrum of (*E*)-4-(4-(Dimethylamino)benzylidene)-3-methyl-1-phenyl-1*H*-pyrazol-5(4*H*)-one.




Figure 24c: FT-IR (KBr) spectrum of (*E*)-4-(4-(Dimethylamino)benzylidene)-3-methyl-1-phenyl-1*H*-pyrazol-5(4*H*)-one.