

Supplementary Material

Selective decolorization of cationic dyes by peroxymonosulfate: non-radical mechanism and effect of chloride

Yang Lei ^{a, b}, Chuh-Shun Chen ^b, Jia Ai^a, Heng Lin^a, Yao-Hui Huang ^{b, c*}, Hui Zhang

^{a*}

^a Department of Environmental Engineering, Wuhan University, Wuhan.

^b Department of Chemical Engineering, National Cheng Kung University, Tainan.

^c Sustainable Environment Research Center, National Cheng Kung University, Tainan.

*Corresponding author

Phone: 86-27-68775837;

Fax: 86-27-68778893;

E-mail: eeng@whu.edu.cn. (H. Zhang)

Phone: 886-2757575x62636;

Fax: 886-6-2344496;

E-mail: yhhuang@mail.ncku.edu.tw. (Y.H. Huang)




Figure S1. Influence of $ZnFe_2O_4$ dosage on the degradation of RhB in PMS/ $ZnFe_2O_4$ system. Conditions: $[PMS] = 2.5$ mM, $[RhB] = 10$ mg/L, $pH_0 = 4.5 \pm 0.1$.




Figure S2. Influence of Ar purging on the degradation of RhB. Conditions: [PMS] = 2.5 mM, $\text{pH}_0 = 4.5 \pm 0.1$. [Ar] = 3 L/min, [RhB] = 10 mg/L.

Table S1. The final pH of RhB solution at different initial solution pH. Conditions: [PMS] = 2.5 mM, [RhB] = 10 mg/L, Time = 120 min.

Initial pH	2.0	4.5	7.0	10.0	12.0
After adding PMS	1.92	2.57	2.56	2.55	11.72
Final pH	0.97	2.53	2.5	2.46	11.56