

Supporting Information Section:

Role of Tryptophan 135 of Chandipura Virus Phosphoprotein P in Dimerization and Complex Formation with Leader RNA: Structural Aspect using Time Resolved Anisotropy and Simulation

Manini Mukherjee^a, Aditya Sarkar^b, Arunava Roy^c, Pinki Saha Sardar^a, Ansuman Lahiri^{b*}, Dhruvajyoti Chattopadhyay^{c*} and Sanjib Ghosh^{a*}

^aDepartment of Chemistry, Presidency University, Kolkata 700 073, India.

^bDepartment of Bio-Physics, Molecular Biology and Bioinformatics, University of Calcutta, Kolkata - 700 009, India.

^cDr. B. C. Guha Centre for Generic Engineering and Biotechnology, Department of Biotechnology, University of Calcutta, Kolkata 700019, India.

***Author for correspondence:**

Tel.: +91-9836940620 (S.G.), +9133-2241 3893 (S.G.); Fax: +9133-2241 3893 (S.G.);

E-mail: sanjibg@cal2.vsnl.net.in (S.G.); pchemsg@gmail.com (S.G.)




Figure S1: Steady state fluorescence spectra at 298 K of WT CHPV and the mutants W135F and W105F in 40% EG- buffer at (A) 300 nM concentration and (B) 1500 nM concentration. $\lambda_{\text{exc}} = 295$ nm, Excitation and emission bandpass = 10 nm and 5 nm each.