

SUPPLEMENTARY INFORMATION

Unravelling the onset of the exchange bias effect in Ni(core)@NiO(shell) nanoparticles embedded in a mesoporous carbon matrix

Natalia Rinaldi-Montes,^{*,a} Pedro Gorria,^b David Martínez-Blanco,^c Zakariae Amghouz,^c Antonio B. Fuertes,^d Luis Fernández Barquín,^e Imanol de Pedro,^e Luca Olivi,^f and Jesús A. Blanco^a

^a Departamento de Física, Universidad de Oviedo, E-33007 Oviedo, Spain.

^b Departamento de Física & IUTA, EPI, Universidad de Oviedo, E-33203 Gijón, Spain.

^c Servicios Científico-Técnicos, Universidad de Oviedo, E-33006 Oviedo, Spain.

^d Instituto Nacional del Carbón (CSIC), E-33080 Oviedo, Spain.

^e *CITIMAC*, Facultad de Ciencias, Universidad de Cantabria, E-39005 Santander, Spain.

^f Elettra-Sincrotrone Trieste S.C.p.A., 34149 Basovizza, Trieste, Italy.

Corresponding author E-mail: nataliarin@gmail.com

1. Size distribution of the nanoparticles




Figure S1. (Color online) Histograms of the particle size distributions of the samples together with log-normal fits, providing mean NP diameters (D_{TEM}) and standard deviations (σ).

2. $M(H)$ curves measured at $T = 300$ K




Figure S2. (Color online) (Left) $M(H)$ curves for samples T673, T773 and T873 (empty circles) measured at room temperature ($T = 300$ K). Lines represent the best fit of the experimental data to a combination of the Langevin function and the lognormal size distribution.⁶⁰ (Right) Enlarged views of the central part of the left $M(H)$ curves.




Figure S3. (Color online) (Left) $M(H)$ curves for samples T1023 and T1173 (empty circles) measured at room temperature ($T = 300 \text{ K}$), showing a small hysteresis loop. Lines provide guides for the eyes. (Right) Enlarged views of the central part of the left $M(H)$ curves.