

Titania supported MOF-199 derived Cu-Cu₂O nanoparticles: Highly efficient non noble metal photocatalysts for hydrogen production from alcohol-water mixtures

Received 00th January 20xx,
Accepted 00th January 20xx

DOI: 10.1039/x0xx00000x

www.rsc.org/

Imran Majeed,¹ Muhammad Amtiaz Nadeem,^{2**} Amin Badshah,¹ Fehmida Kanodarwala,³ Hassan Ali,¹ M. Abdullah Khan,⁴ John Arron Stride,³ and Muhammad Arif Nadeem^{1*}




Figure S1: TEM images of TiO₂-MOF-500 °C




Figure S2. XPS core level spectra of Ti 2p, O 1s, C 1s and Cu 2p for TiO₂-MOF-500 °C




Figure S3: H₂ production rates over TiO₂-MOF-400 °C from different concentrations of glycerol

*Catalysis and Nanomaterials Lab 27, Department of Chemistry, Quaid-i-Azam University, Islamabad 45320, Pakistan. manadeem@qau.edu.pk, +92-51-9064-2062

^bDepartment of Environmental Sciences, Quaid-i-Azam University, Islamabad 45320, Pakistan. manadeempk@gmail.com, NadeemMI@SABIC.com

^cSchool of Chemistry, University of New South Wales, Sydney, NSW 2052, Australia.

^dUS. Pakistan Center for Advanced Studies in Energy, NUST, Islamabad, 44000, Pakistan
† Present Address: SABIC-CRD at KAUST.

Electronic Supplementary Information (ESI) available: [details of any supplementary information available should be included here]. See DOI: 10.1039/x0xx00000x