

Supporting Information for

Selective and tunable synthesis of 3-arylsuccinimides and 3-arylmaleimides from arenediazonium tetrafluoroborates and maleimides

Zhen-Hua Yang^{a,1}, Zhong-Hui Chen^{a,1}, Yu-Long An^a, and Sheng-Yin Zhao^{*a,b}

^aDepartment of Chemistry, Donghua University, No.2999 North Renmin Road,
Shanghai 201620, P. R. of China;

^bState Key Laboratory of Bioorganic & Natural Products Chemistry, Shanghai Institute of
Organic Chemistry, Chinese Academy of Sciences, Shanghai 200032, P. R. of China.

Tel: +86 21 67792436; Fax: +86 21 67792608-805;

E-mail: syzhao8@dhu.edu.cn

Content:


¹H NMR and ¹³C NMR spectra of the products.

¹H NMR and ¹³C NMR spectra


¹H and ¹³C NMR spectra were recorded using a Bruker AV 400 MHz spectrometer in CDCl₃ or DMSO-d₆ with tetramethylsilane as internal standard.

3-phenylpyrrolidine-2,5-dione (4a) in CDCl₃

¹H NMR




¹³C NMR




3-(4-nitrophenyl)-pyrrolidine-2,5-dione (4b) in CDCl_3

^1H NMR




^{13}C NMR




3-(4-chlorophenyl)-pyrrolidine-2,5-dione (4c) in CDCl₃

¹H NMR




¹³C NMR




3-(4-bromophenyl)-pyrrolidine-2,5-dione (4d) in CDCl₃

¹H NMR




¹³C NMR




3-(4-methoxyphenyl)-pyrrolidine-2,5-dione (4e) in CDCl_3

^1H NMR




^{13}C NMR




3-(2-chloro-6-methylphenyl)-pyrrolidine-2,5-dione (4f) in CDCl₃

¹H NMR




¹³C NMR




3-(Naphthalen-2-yl)-pyrrolidine-2,5-dione (4g) in DMSO-d₆

¹H NMR




¹³C NMR




1-methyl-3-phenyl-pyrrolidine-2,5-dione (4h) in CDCl₃

¹H NMR




¹³C NMR




3-(4-nitrophenyl)-1-phenyl-Pyrrolidine-2,5-dione (4i) in CDCl_3


^1H NMR




^{13}C NMR



3-(3-chloro-4-trifluoromethylphenyl)-1-phenyl-Pyrrolidine-2,5-dione (4j) in CDCl₃




¹³C NMR




3-(4-methoxyphenyl)-1-phenyl-pyrrolidine-2,5-dione (4k) in DMSO-d₆

¹H NMR




¹³C NMR




3-(4-methoxyphenyl)-1-benzyl-pyrrolidine-2,5-dione (4l) in DMSO-d₆

¹H NMR




¹³C NMR




3-phenyl-1H-pyrrole-2,5-dione (5a**) in DMSO-d₆**

¹H NMR




¹³C NMR




3-(3-chloro-4-trifluoromethylphenyl)-1H--pyrrole-2,5-dione (5b**) in DMSO-d₆**

¹H NMR




¹³C NMR




3-(4-bromophenyl)-1H-pyrrole-2,5-dione (5c) in DMSO-d₆

¹H NMR




¹³C NMR




3-(4-chlorophenyl)-1H-pyrrole-2,5-dione (5d) in DMSO-d₆

¹H NMR




¹³C NMR




3-(4-methoxyphenyl)-1H-pyrrole-2,5-dione (5e**) in CDCl₃**

¹H NMR




¹³C NMR




3-(4-bromophenyl)-1-phenyl-1H-pyrrole-2,5-dione (5f**) in DMSO-d₆**

¹H NMR




¹³C NMR



3-(4-nitrophenyl)-1H-pyrrole-2,5-dione (5g) in DMSO

^1H NMR



^{13}C NMR

