RSC Advances

Hydrophilic ionic liquid-assisted control of size and morphology of ZnO nanoparticles prepared by chemical precipitation method[†]

Mousumi Akter,^a Shazia Sharmin Satter,^a Ajaya Kumar Singh,^b M. Muhibur Rahman,^c M.

Yousuf A. Mollah^c, and Md. Abu Bin Hasan Susan^{a*}

^aDepartment of Chemistry, University of Dhaka, Dhaka 1000, Bangladesh

^bDepartment of Chemistry, Govt. V. Y. T. PG. Autonomous College, Chhattisagarh, India

^cUniversity Grants Commission of Bangladesh, 29/1 Agargaon, Sher-E-Banla Naar, Dhaka

1207, Bangladesh

[†]Electronic supplementary information (ESI)

Fig. S1 EDX spectrum of ZnO NPs.

- Fig. S2 Stucture of [EMIM][MeSO₄]
- Fig. S3 Zeta potential distribution curve of ZnO NPs in the absence of ionic liquid.

Fig. S4 Zeta potential distribution curve of ZnO NPs in 5.0% IL(v/v).

Fig. S5 Size distribution curve for [EMIM][MeSO₄].

Fig. S6 SEM images (filament technique) of ZnO NPs synthesized in absence of [EMIM][MeSO₄] (a) and in presence of (b) 0.1% (c) 1.5% (d) 2% (e) 3.5% (f) 5% (v/v) [EMIM][MeSO₄]




Fig. S1 EDX spectrum of ZnO NPs.




Fig. S2 Stucture of [EMIM][MeSO₄].




Fig. S3 Zeta potential distribution curve of ZnO NPs in absence of ionic liquid.




Fig. S4 Zeta potential distribution curve of ZnO NPs in 5% IL(v/v).




Fig. S5 Size distribution curve for [EMIM][MeSO₄].




Fig. S6 SEM images of ZnO NPs synthesized in absence of [EMIM][MeSO4] (a) and in presence of (b) 0.1% (c) 1.5% (d) 2% (e) 3.5% (f) 5% (v/v) [EMIM][MeSO4].