

Electronic supplementary information (ESI)

Contents

Fig. S1 TEM image of g-C ₃ N ₄	1
Fig. S2 Time resolved photoluminescence spectra of g-C ₃ N ₄ and HCNNSs excited at 337 nm and monitored at emission 500 nm.	2
Fig. S3 Ion chromatographic effluent curve for LEVO photocatalytic degradation products over visible-light irradiating HCNNS-200. Catalyst amount 50 mg; $c_0 = 20$ mg L ⁻¹ ; volume 50 mL; $400\text{ nm} < \lambda < 680\text{ nm}$	3
Fig. S4 ESI-MS identified intermediate products yielded during the process of visible-light photocatalytic degradation of LEVO. $400\text{ nm} < \lambda < 680\text{ nm}$; catalyst amount 50 mg; $C_0 = 20\text{ mg L}^{-1}$, volume 50 mL.	4

Fig. S1 TEM image of g-C₃N₄.




Fig. S2 Time resolved photoluminescence spectra of g-C₃N₄ and HCNNSs excited at 337 nm and monitored at emission 500 nm.




Fig. S3 Ion chromatographic effluent curve for LEVO photocatalytic degradation products over visible-light irradiating HCNNS-200. Catalyst amount 50 mg; $c_0 = 20$ mg L⁻¹; volume 50 mL; 400 nm < λ < 680 nm.




Fig. S4 ESI-MS identified intermediate products yielded during the process of visible-light photocatalytic degradation of LEVO. $400\text{ nm} < \lambda < 680\text{ nm}$; catalyst amount 50 mg; $C_0 = 20\text{ mg L}^{-1}$, volume 50 mL.

