

Electronic supplementary information

PEG-b-polydisulfide-L-lysine based redox responsive cationic polymer for efficient gene transfection

Ihsan Ullah^{a#}, Jing Zhao^{a,b#}, Shah Rukh^c, Khan Muhammad^a, Jintang Guo^{a,b}, Xiangkui Ren^{*a,b,d}, Shihai Xia^{*e}, Wencheng Zhang^f, Yakai Feng^{*a,b,d}

a. School of Chemical Engineering and Technology, Tianjin University, Yaguan Road 135, Tianjin 300350, China

b. Collaborative Innovation Centre of Chemical Science and Chemical Engineering (Tianjin), Weijin Road 92, Tianjin 300072, China

c. Department of Chemistry, School of Science Abdul Wali Khan University, Mardan 23200 Pakistan.

d. Key Laboratory of Systems Bioengineering of Ministry of Education, Tianjin University, Weijin Road 92, Tianjin 300072, China.

e. Department of Hepatopancreatobiliary and Splenic Medicine, Affiliated Hospital, Logistics University of People's Armed Police Force, 220 Chenglin Road, Tianjin 300162, China.

f. Department of Physiology and Pathophysiology, Logistics University of Chinese People's Armed Police Force, Tianjin 300309, China

g.

#Ihsan Ullah and Jing Zhao contributed equally to this work.

*Corresponding Author: Y. Feng, School of Chemical Engineering and Technology, Tianjin University, Tianjin 300350, China. E-mail: yakai.feng@tju.edu.cn




Figure S1 ^1H NMR spectra of PEG(OH) $_4$ (A) and PEG(Ac) $_4$ (B).