

Supporting Information

Palladium complexes of N-heterocyclic carbenes displaying an unsymmetrical N-alkylfluorenyl/N'-aryl substitution pattern and their behaviour in Suzuki-Miyaura cross coupling

Hamzé Almallah,^{a,b} Eric Brenner,^{*a} Dominique Matt,^{*a} Jack Harrowfield,^c Mohamad Jahjah^b and Akram Hijazi^b

^a*Laboratoire de Chimie Inorganique Moléculaire et Catalyse, Institut de Chimie (UMR 7177 CNRS), Université de Strasbourg, 4 rue Blaise Pascal, 67070 Strasbourg Cedex, France. E-mail: eric.brenner@unistra.fr, dmatt@unistra.fr*

^b*Inorganic and Organometallic Coordination Chemistry Laboratory, Faculty of Sciences, Lebanese University, R. Hariri University Campus, Beyrouth, Hadath, Lebanon*


^c*Institut de Science et Ingénierie Supramoléculaires (ISIS), UMR 7606 CNRS, Université de Strasbourg, 8 rue Gaspard Monge, 67083 Strasbourg cedex, France*

Contents:


1) ^1H , $^{13}\text{C}\{^1\text{H}\}$ and ^{19}F NMR spectra ————— S1-S30

1) ^1H , $^{13}\text{C}\{^1\text{H}\}$ and ^{19}F NMR spectra


^1H NMR (CDCl_3 , 300 MHz) of imidazolium 3^H




$^{13}\text{C}\{^1\text{H}\}$ and DEPT NMR (CDCl_3 , 125 MHz) of imidazolium 3^H




¹H NMR ([D₆]DMSO, 500 MHz) of imidazolium **4^H**




¹³C{¹H} and DEPT NMR ([D₆]DMSO, 125 MHz) of imidazolium **4^H**




^1H NMR (CDCl_3 , 500 MHz) of imidazolium $3^{*\text{Et}}$




$^{13}\text{C}\{^1\text{H}\}$ and DEPT NMR (CDCl_3 , 125 MHz) of imidazolium $3^{*\text{Et}}$




¹H NMR (CDCl_3 , 500 MHz) of imidazolium **3^{Et}**




¹³C{¹H} and DEPT NMR (CDCl_3 , 125 MHz) of imidazolium **3^{Et}**




¹⁹F NMR (CDCl_3 , 282 MHz) of imidazolium **3^{Et}**




¹H NMR (CDCl_3 , 500 MHz) of imidazolium **4*Et**




$^{13}\text{C}\{\text{H}\}$ and DEPT NMR (CDCl_3 , 125 MHz) of imidazolium $\mathbf{4}^*\text{Et}$




^1H NMR (CDCl_3 , 500 MHz) of imidazolium $\mathbf{4}\text{Et}$




$^{13}\text{C}\{\text{H}\}$ and DEPT NMR (CDCl_3 , 125 MHz) of imidazolium **4^{Et}**




^{19}F NMR (CDCl_3 , 282 MHz) of imidazolium **4^{Et}**




^1H NMR (CDCl_3 , 500 MHz) of imidazolium $3^{*\text{Pr}}$




$^{13}\text{C}\{^1\text{H}\}$ and DEPT NMR (CDCl_3 , 125 MHz) of imidazolium $3^{*\text{Pr}}$




¹H NMR (CDCl_3 , 500 MHz) of imidazolium **3^{Pr}**




¹³C{¹H} and DEPT NMR (CDCl_3 , 125 MHz) of imidazolium **3^{Pr}**




¹⁹F NMR (CDCl_3 , 282 MHz) of imidazolium **3^{Pr}**




¹H NMR (CDCl_3 , 400 MHz) of imidazolium **4*Pr**




$^{13}\text{C}\{\text{H}\}$ and DEPT NMR (CDCl_3 , 125 MHz) of imidazolium $\mathbf{4}^*\text{Pr}$




^1H NMR (CDCl_3 , 500 MHz) of imidazolium $\mathbf{4}^*\text{Pr}$




$^{13}\text{C}\{\text{H}\}$ and DEPT NMR (CDCl_3 , 125 MHz) of imidazolium **4^{Pr}**




^{19}F NMR (CDCl_3 , 282 MHz) of imidazolium **4^{Pr}**




¹H NMR (CDCl_3 , 500 MHz) of imidazolium **3^{EOM}**




¹³C{¹H} and DEPT NMR (CDCl_3 , 125 MHz) of imidazolium **3^{EOM}**




¹⁹F NMR (CDCl_3 , 282 MHz) of imidazolium **3^{EOM}**




¹H NMR (CDCl_3 , 500 MHz) of imidazolium **4^{EOM}**





$^{13}\text{C}\{\text{H}\}$ and DEPT NMR (CDCl_3 , 125 MHz) of imidazolium **4^{EOM}**




^{19}F NMR (CDCl_3 , 282 MHz) of imidazolium **4^{EOM}**




^1H NMR (CDCl_3 , 500 MHz) of complex $\mathbf{5}^{\text{Et}}$




¹H NMR (CDCl_3 , 500 MHz) of complex **6^{Et}**




¹³C{¹H} and DEPT NMR (CDCl_3 , 125 MHz) of complex **6^{Et}**




¹H NMR (CDCl_3 , 500 MHz) of complex **5^{Pr}**




¹³C{¹H} and DEPT NMR (CDCl_3 , 125 MHz) of complex **5^{Pr}**




¹H NMR (CDCl_3 , 400 MHz) of complex **6^{Pr}**




¹³C{¹H} and DEPT NMR (CDCl_3 , 125 MHz) of complex **6^{Pr}**




¹H NMR (CDCl_3 , 400 MHz) of complex **5^{EOM}**




¹³C{¹H} and DEPT NMR (CDCl_3 , 125 MHz) of complex **5^{EOM}**




¹H NMR (CDCl_3 , 500 MHz) of complex **6^{EOM}**




¹³C{¹H} and DEPT NMR (CDCl_3 , 125 MHz) of complex **6^{EOM}**




^1H NMR ($[\text{D}_6]\text{DMSO}$, 500 MHz) of imidazolium 7^H




$^{13}\text{C}\{\text{H}\}$ and DEPT NMR ($[\text{D}_6]\text{DMSO}$, 125 MHz) of imidazolium 7^H




¹⁹F NMR ([D₆]DMSO, 282 MHz) of imidazolium **7^H**




¹H NMR (CDCl₃, 400 MHz) of imidazolium **8^{Me}**




$^{13}\text{C}\{\text{H}\}$ and DEPT NMR (CDCl_3 , 125 MHz) of imidazolium **8^{Me}**




^{19}F NMR (CDCl_3 , 470 MHz) of imidazolium **8^{Me}**





¹H NMR (CDCl_3 , 500 MHz) of imidazolium **8^{Et}**




¹³C{¹H} and DEPT NMR (CDCl_3 , 125 MHz) of imidazolium **8^{Et}**




¹⁹F NMR (CDCl_3 , 282 MHz) of imidazolium **8^{Et}**




¹H NMR (CDCl_3 , 500 MHz) of imidazolium **8^{Pr}**




$^{13}\text{C}\{\text{H}\}$ and DEPT NMR (CDCl_3 , 125 MHz) of imidazolium **8^{Pr}**




^{19}F NMR (CDCl_3 , 282 MHz) of imidazolium **8^{Pr}**




¹H NMR (CDCl_3 , 500 MHz) of complex **9^{Me}**




¹³C{¹H} and DEPT NMR (CDCl_3 , 125 MHz) of complex **9^{Me}**



¹H NMR (CDCl_3 , 400 MHz) of complex **9^{Et}**



¹H NMR (CDCl_3 , 500 MHz) of complex **9^{Pr}**



$^{13}\text{C}\{\text{H}\}$ and DEPT NMR (CDCl_3 , 125 MHz) of complex **9^{Pr}**

