

Pd-Catalyzed Sequential Hydroarylation and Olefination Reactions of 3-AllylChromones

Yi-En Liang, Chia-Ling Lu, and Wen-Tai Li*

National Research Institute of Chinese Medicine, Ministry of Health and Welfare,
Taipei 11221, Taiwan, R.O.C.

*Email: wtli@nricm.edu.tw


Supporting Information

| Table of Contents | Page No |
|---|---------|
| 1. ^1H and ^{13}C NMR spectra | S2 |
| 2. ORTEP diagram of 2c | S63 |
| 3. X-ray data of 2c | S64 |

3-Allyl-7-methoxy-4H-chromen-4-one (1a)

¹H NMR


PROTON_01 — J45_CDCl₃_600 2018/02/01 —



3-Allyl-4H-chromen-4-one (1b)




¹H NMR

PROTON_01 — J47_CDCl₃_600 2018/02/01 —





*3-Allyl-6-chloro-4H-chromen-4-one (**1d**)*

¹H NMR





(E)-3-(But-2-en-1-yl)-4H-chromen-4-one (3a)

¹H NMR




¹³C NMR




3-(But-2-en-1-yl)-7-methoxy-4H-chromen-4-one (3b)

¹H NMR




¹³C NMR




3-(But-2-en-1-yl)-6-chloro-4H-chromen-4-one (3c)

¹H NMR




¹³C NMR




3-Cinnamyl-7-methoxy-4H-chromen-4-one (5a)

¹H NMR





¹³C NMR




*(E)-3-(3-(*p*-Tolyl)allyl)-4*H*-chromen-4-one (**5b**)*

¹H NMR




*(E)-7-Methoxy-3-(3-(*p*-tolyl)allyl)-4*H*-chromen-4-one (5c)*

¹H NMR




¹³C NMR




(E)-7-(3-Phenylprop-1-en-1-yl)-8H-[1,3]dioxolo[4,5-g]chromen-8-one (5f)

¹H NMR




¹³C NMR




*7-Methoxy-3-(1-(*p*-tolyl)propyl)-4*H*-chromen-4-one (2a)*

¹H NMR



¹³C NMR




*3-(1-(*o*-Tolyl)propyl)-4*H*-chromen-4-one (2b')*

¹H NMR

PROTON_01 — J232_A12-14_A8-12_CDCl₃_500 2019/01/15 —


8.212 7.820 7.349 7.337 7.277 7.24 7.23 7.191 7.176 7.13 7.159 7.145 7.121 7.119 7.107 7.104 4.401 4.389 4.383 4.371 2.36 2.049 2.037 2.022 2.010 2.007 1.995 1.908 1.904 1.934 1.934 1.90 1.80 1.776 1.666 1.662 1.648 0.959 0.944 0.929



¹³C NMR


CARBON_01 — J232_A12-14_A8-12_CDCl₃_500 2019/01/15 —

177.063 156.135 153.263 149.295 140.224 137.332 133.332 130.710 128.148 126.271 126.217 126.090 125.879 124.860 123.894 117.947 77.254 77.090 76.745 37.929 27.280 19.792 12.534




3-(1-(2,4-Dimethylphenyl)propyl)-7-methoxy-4H-chromen-4-one (2c)

¹H NMR




¹³C NMR




3-(1-(4-Hydroxyphenyl)propyl)-7-methoxy-4H-chromen-4-one (2e).

¹H NMR




¹³C NMR




3-(1-(2-Hydroxyphenyl)propyl)-7-methoxy-4H-chromen-4-one (2e').

¹H NMR




¹³C NMR




7-Methoxy-3-(1-(naphthalen-1-yl)propyl)-4H-chromen-4-one (2f)

¹H NMR




¹³C NMR




7-Methoxy-3-(1-(naphthalen-2-yl)propyl)-4H-chromen-4-one (2f')


¹H NMR




¹³C NMR



7-Methoxy-3-(1-(4-phenoxyphenyl)propyl)-4H-chromen-4-one (2g)
¹H NMR




¹³C NMR




7-Methoxy-3-(1-(4-oxochroman-6-yl)propyl)-4H-chromen-4-one (2i)

¹H NMR




¹³C NMR




3-(1-Mesilylpropyl)-7-methoxy-4H-chromen-4-one (2j)

¹H NMR




¹³C NMR




3-(1-(4-Acetyl-3-hydroxyphenyl)propyl)-7-methoxy-4H-chromen-4-one (2l)

¹H NMR




¹³C NMR




3-(1-(3-Acetyl-2-hydroxyphenyl)propyl)-7-methoxy-4H-chromen-4-one (2l')

¹H NMR





¹³C NMR




4-(7-Methoxy-4-oxo-4H-chromen-3-yl)-4-(*p*-tolyl)butanoic acid (2m**)**

¹H NMR




3-(1-(*p*-Tolyl)butyl)-4*H*-chromen-4-one (4a**)**

¹H NMR




¹³C NMR




7-Methoxy-3-(1-(2,4-dimethylphenyl)butyl)-4H-chromen-4-one (4b)

¹H NMR




¹³C NMR




6-Chloro-3-(1-(2,5-dimethylphenyl)butyl)-4H-chromen-4-one (4c**)**

¹H NMR




¹³C NMR



3-(1-(3,4-Dimethylphenyl)butyl)-4H-chromen-4-one (4d)


¹H NMR

PROTON_01 — J211_A6-7_A37-43_CDCl₃_500 2019/01/19 —




¹³C NMR

CARBON_01 — J211_A6-7_A37-43_CDCl₃_500 2019/01/19 —




*7-Methoxy-3-(3-phenyl-1-(*p*-tolyl)propyl)-4*H*-chromen-4-one (**6a**)*

¹H NMR




¹³C NMR




3-(1,3-Di-p-tolylpropyl)-4H-chromen-4-one (6b**)**

¹H NMR




¹³C NMR




3-(1,3-Di-p-tolylpropyl)-7-methoxy-4H-chromen-4-one (6c)

¹H NMR





¹³C NMR



*3-(1-(3,4-Dimethylphenyl)-3-(*p*-tolyl)propyl)-4*H*-chromen-4-one (**6d**)*


¹H NMR

PROTON_01 — J215_A3-10_A22-28_CDCl₃_500 2019/01/19 —




¹³C NMR

CARBON_01 — J215_A3-10_A22-28_CDCl₃_500 2019/01/19 —




3-(1-(2,5-Dimethylphenyl)-3-phenylpropyl)-7-methoxy-4H-chromen-4-one (6e)

¹H NMR




¹³C NMR




*7-(1-(4-Methoxyphenyl)-3-phenylpropyl)-8*H*-[1,3]dioxolo[4,5-*g*]chromen-8-one (**6f**)*

¹H NMR




¹³C NMR




*7-(1-(2-Methoxyphenyl)-3-phenylpropyl)-8*H*-[1,3]dioxolo[4,5-*g*]chromen-8-one (**6f'**)*

¹H NMR




¹³C NMR




6-Chloro-3-(1,3-di-p-tolylpropyl)-4H-chromen-4-one (6g)

¹H NMR




¹³C NMR




3-(1-([1,1'-Biphenyl]-4-yl)-3-(*p*-tolyl)propyl)-7-methoxy-4H-chromen-4-one (*6h*)

¹H NMR




¹³C NMR




7-Methoxy-3-(3-(2,4-dimethylphenyl)-3-phenylpropyl)-4H-chromen-4-one (6i)

¹H NMR




¹³C NMR




Butyl (E)-3-(3,5-dimethyl-2-(1-(4-oxo-4H-chromen-3-yl)propyl)phenyl)acrylate (7a)

¹H NMR




¹³C NMR




Methyl (E)-3-(3,5-dimethyl-2-(1-(4-oxo-4H-chromen-3-yl)propyl)phenyl)acrylate (7b)

¹H NMR




¹³C NMR




Butyl (E)-3-(2-(1-(6-fluoro-4-oxo-4H-chromen-3-yl)propyl)-3,5-dimethylphenyl)acrylate (7c)

¹H NMR




¹³C NMR




Ethyl (E)-3-(2-(1-(7-methoxy-4-oxo-4H-chromen-3-yl)propyl)-5-methylphenyl)acrylate (7d)

¹H NMR




¹³C NMR




Ethyl (E)-3-(5-methyl-2-(1-(4-oxo-4H-chromen-3-yl)propyl)phenyl)acrylate (7e)


¹H NMR




¹³C NMR



Ethyl (E)-3-(3-methyl-2-(1-(4-oxo-4H-chromen-3-yl)propyl)phenyl)acrylate (7e')
¹H NMR




¹³C NMR




Diethyl 3,3'-(5-methyl-2-(1-(4-oxo-4H-chromen-3-yl)propyl)-1,3-phenylene)(2E,2'E)-diacrylate (7e")

¹H NMR




¹³C NMR




Butyl (E)-3-(2-(1-(7-methoxy-4-oxo-4H-chromen-3-yl)propyl)-4,5-dimethylphenyl)acrylate (7f)

¹H NMR




¹³C NMR




Butyl (E)-3-(2-(1-(7-methoxy-4-oxo-4H-chromen-3-yl)propyl)-5-methylphenyl)acrylate (7g)

¹H NMR




¹³C NMR




Butyl(E)-3-(2-(1-(7-methoxy-4-oxo-4H-chromen-3-yl)propyl)-3methylphenyl)acrylate
(7g')

¹H NMR




¹³C NMR




Dibutyl 3,3'-(2-(1-(7-methoxy-4-oxo-4H-chromen-3-yl)propyl)-5-methyl-1,3-phenylene)(2E,2'E)-diacrylate (7g")

¹H NMR




¹³C NMR




(E)-3-(1-(2,4-Dimethyl-6-(2-(phenylsulfonyl)vinyl)phenyl)propyl)-4H-chromen-4-one

(7h)

¹H NMR





¹³C NMR



Ethyl (E)-3-(3,5-dimethyl-2-(1-(4-oxo-4H-chromen-3-yl)propyl)phenyl)acrylate (7i)


¹H NMR




Butyl (E)-3-(2-(1-(6-fluoro-4-oxo-4H-chromen-3-yl)propyl)-5-methylphenyl)acrylate

(7j)

¹H NMR





¹³C NMR




Butyl (E)-3-(2-(1-(6-fluoro-4-oxo-4H-chromen-3-yl)propyl)-3-methylphenyl)acrylate
(7j)
¹H NMR

PROTON_01 — J261_A9-11_A38-43_CDCl₃_500 2019/01/22 — —




¹³C NMR




Dibutyl 3,3'-(2-(1-(6-fluoro-4-oxo-4H-chromen-3-yl)propyl)-5-methyl-1,3-phenylene)(2E,2'E)-diacrylate (7j")

¹H NMR




¹³C NMR




Butyl (E)-3-(5-methyl-2-(1-(4-oxo-4H-chromen-3-yl)butyl)phenyl)acrylate (7k)

¹H NMR




¹³C NMR




Methyl (E)-3-(5-methyl-2-(1-(4-oxo-4H-chromen-3-yl)butyl)phenyl)acrylate (7l)


¹H NMR




¹³C NMR



Methyl (E)-3-(3-methyl-2-(1-(4-oxo-4H-chromen-3-yl)butyl)phenyl)acrylate (7l')
¹H NMR




¹³C NMR




Methyl (E)-3-(5-methyl-2-(1-(4-oxo-4H-chromen-3-yl)butyl)phenyl)acrylate (7l")

¹H NMR




¹³C NMR



Single crystal X-ray diffraction

The single crystal was obtained by slow evaporation of **2c** in a solution of acetone and n-hexane. The structure was solved with the Oxford Gemini dual system dingle-crystal XRD equipped with Cryojet, and refined with the SHELXL-2014 using Least Squares minimization. The X-ray crystallographic data can be obtained free of charge from the Cambridge Crystallographic Data Center via www.ccdc.cam.ac.uk/data_request/cif.



IC19207 in P-1




Figure S1. ORTEP plot of the crystallographic structure of **2c**. Thermal probability ellipsoids are shown at the 50% probability level.

Table S1. Crystal data and experimental details for ic19207 (**2c**).

| Crystal data | | |
|-----------------------------------|---|--|
| Empirical formula | C21 H22 O3 | |
| Formula weight | 322.38 | |
| Crystal system | Triclinic | |
| Space group | P-1 | |
| Unit cell dimensions | a = 6.3862(3) Å b = 8.2346(4) Å c = 17.2255(12) Å | α = 100.358(5)°. β = 97.648(5)°. γ = 95.824(4)°. |
| Volume | 875.83(9) Å ³ | |
| Z | 2 | |
| F(000) | 344 | |
| Density (calculated) | 1.222 Mg/m ³ | |
| Wavelength | 0.71073 Å | |
| Cell parameters reflections used | 2406 | |
| Theta range for Cell parameters | 4.0070 to 28.7000°. | |
| Absorption coefficient | 0.081 mm ⁻¹ | |
| Temperature | 150(2) K | |
| Crystal size | 0.20 x 0.20 x 0.15 mm ³ | |
| Data collection | | |
| Diffractometer | Xcalibur, Atlas, Gemini | |
| Absorption correction | Semi-empirical from equivalents | |
| Max. and min. transmission | 1.00000 and 0.95886 | |
| No. of measured reflections | 5933 | |
| No. of independent reflections | 3771 [R(int) = 0.0194] | |
| No. of observed [I>2_igma(I)] | 2764 | |
| Completeness to theta = 25.242° | 99.2 % | |
| Theta range for data collection | 3.152 to 27.497°. | |
| Refinement | | |
| Final R indices [I>2sigma(I)] | R1 = 0.0618, wR2 = 0.1916 | |
| R indices (all data) | R1 = 0.0861, wR2 = 0.2156 | |
| Goodness-of-fit on F ² | 1.329 | |
| No. of reflections | 3771 | |
| No. of parameters | 217 | |
| No. of restraints | 0 | |
| Largest diff. peak and hole | 0.414 and -0.407 e.Å ⁻³ | |