

## Supplementary Data

### Performance of selenate removal by biochar embedded nano zero-valent iron and the biological toxicity to *Escherichia coli*

Liping Liang<sup>a,b\*</sup>, Yuanyuan Xue<sup>a</sup>, Gangliang Tian<sup>b</sup>, Qiaole Mao<sup>b</sup>, Zixuan Lou<sup>b</sup>, Qian Wu<sup>a</sup>, Qian Wang<sup>a</sup>, Juanshan Du<sup>c</sup>, and Xu Meng<sup>d\*</sup>

a. School of Civil Engineering, Shaoxing University, Shaoxing 312000, P. R. China

b. College of Life Science, Shaoxing University, Shaoxing 312000, P. R. China

c. State Key Laboratory of Urban Water Resource and Environment, Harbin Institute of Technology, Harbin 150090, PR China.

d. College of Textile and Garment, Shaoxing University, Shaoxing 312000, P. R. China

\*Corresponding author contact information:

Liping Liang, email: Liangliping0702@163.com; Tel.: +86-575-88345007.

Xu Meng, email: mengxu0@163.com; Tel.: +86-575-88341506.

Number of pages (including this page): 4

Number of Figures: 2

Number of Tables: 1




Fig. S1 Particle size distribution of the biochar sample




Fig. S2 The EDS elemental mapping of (a) all elements, (b) C, (c) Fe, and (d) O elements of the BC-nZVI sample

Table S1 Organic elemental analysis of biochar

N(%)	C(%)	H(%)	S(%)	O(%)
1.091	67.324	3.022	0.059	/