

An original synthesis of highly ordered organosilica with a high content of thiol groups.


Johan Alauzun, Ahmad Mehdi*, Catherine Reyé, Robert J.P. Corriu*

^aLaboratoire de Chimie Moléculaire et Organisation du Solide, UMR 5637 CNRS, Université Montpellier II, Sciences et Techniques du Languedoc, 34095 Montpellier, France.

E-mail : mehdi@univ-montp2.fr

Supplementary Information 1

²⁹Si MAS NMR spectra of X2c.



An original synthesis of highly ordered organosilica with a high content of thiol groups.



Johan Alauzun, Ahmad Mehdi*, Catherine Reyé, Robert J.P. Corriu*

^aLaboratoire de Chimie Moléculaire et Organisation du Solide, UMR 5637 CNRS, Université Montpellier II, Sciences et Techniques du Languedoc, 34095 Montpellier, France.

E-mail : mehdi@univ-montp2.fr

Supplementary Information 2

XRD patterns of X2a and X2b



An original synthesis of highly ordered organosilica with a high content of thiol groups.

Johan Alauzun, Ahmad Mehdi*, Catherine Reyé, Robert J.P. Corriu*

^aLaboratoire de Chimie Moléculaire et Organisation du Solide, UMR 5637 CNRS, Université Montpellier II, Sciences et Techniques du Languedoc, 34095 Montpellier, France.

E-mail : mehdi@univ-montp2.fr

Supplementary Information 3

XRD pattern of **X4c**

