

Electronic Supplementary Information (ESI)

Coenzyme analogs: excellent substitutes (not poor imitations) for electrochemical regeneration

Hye Jung Lee,^a Sahng Ha Lee,^b Chan Beum Park^{*b} and Keehoon Won^{*a}

^a Department of Chemical and Biochemical Engineering, Dongguk University-Seoul, 26 Pil-dong 3-ga, Jung-gu, Seoul 100-715, Republic of Korea. Fax: +82 2 2268 8729; Tel: +82 2 2260 8922; E-mail: keehoon@dongguk.edu

^b Department of Materials Science and Engineering, Korea Advanced Institute of Science and Technology, Science Road 335, Yuseong-gu, Daejeon 305-701, Republic of Korea. Fax: +82 42 350 3310; Tel: +82 42 350 3340; E-mail: parkcb@kaist.ac.kr

Table S1 Absorption maxima (λ_{\max}), molar extinction coefficients (ε), and redox potentials of NAD and its analogs¹¹

NAD and its analogs	λ_{\max}^a (nm)	ε at λ_{\max} (mM⁻¹cm⁻¹)	Redox potential (mV)
NAD	338	6.22	-320
thio-NAD	395	11.30	-285
APAD	363	9.10	-258
PAAD	358	9.30	-262
NAAD	330	5.50	NA

^a The reduced form.