

Electronic Supplementary Information

Synthesis, characterization and staining performance of peripherally and non-peripherally substituted metallo-phthalocyanines bearing 1,3-bis-(trimethylamino)-2-propoxy groups

Mukaddes Özçeşmeci^a, Serap Sancar Baş^b, Barbaros Akkurt^a, Şehnaz Bolkent^{b} and Esin Hamuryudan^{a*}*

^a Istanbul Technical University, Faculty of Science and Letters, Department of Chemistry, 34469 Maslak, Istanbul, Turkey.

^b Istanbul University, Faculty of Science, Department of Biology, 34134 Vezneciler, Istanbul, Turkey

Table of Contents

NMR spectra of 6 and 10	2
Fig. S1 ¹ H NMR spectrum of 6 in d ₆ -DMSO.....	2
Fig. S2 ¹³ C NMR spectrum of 6 in d ₆ -DMSO.....	2
Fig. S3 ¹ H NMR spectrum of 10 in d ₆ -DMSO.....	3
Fig. S4 ¹³ C NMR spectrum of 10 in d ₆ -DMSO.....	3
MALDI-TOF MS spectra of 3 , 5 , 6 , 7 , 9 and 10	4
Fig. S5 Maldi-TOF MS spectrum of 3	4
Fig. S6 Maldi-TOF MS spectrum of 5	4
Fig. S7 Maldi-TOF MS spectrum of 6	5
Fig. S8 Maldi-TOF MS spectrum of 7	5
Fig. S9 Maldi-TOF MS spectrum of 9	6
Fig. S10 Maldi-TOF MS spectrum of 10	6

NMR spectra of 6 and 10




Fig. S1 ^1H NMR spectrum of 6 in d_6 -DMSO.




Fig. S2 ^{13}C NMR spectrum of 6 in d_6 -DMSO.




Fig. S3 ^1H NMR spectrum of **10** in d_6 -DMSO.




Fig. S4 ^{13}C NMR spectrum of **10** in d_6 -DMSO.

MALDI-TOF MS spectra of 3, 5, 6, 7, 9 and 10




Fig. S5 Maldi-TOF MS spectrum of 3.




Fig. S6 Maldi-TOF MS spectrum of 5.




Fig. S7 Maldi-TOF MS spectrum of 6.




Fig. S8 Maldi-TOF MS spectrum of 7.




Fig. S9 Maldi-TOF MS spectrum of **9**.




Fig. S10 Maldi-TOF MS spectrum of **10**.