Supplementary Material

Stöber synthesis of salen-formaldehyde resin polymer- and carbon spheres with high nitrogen content and application of the corresponding Mn-containing carbon spheres as efficient electrocatalysts for the oxygen reduction reaction

Tahereh Hosseinzadeh Sanatkar,^a Alireza Khorshidi,*^a Rouhollah Yaghoubi,^b Esmail Sohouli,^c and Jamaladin Shakeri^d

^a Department of Chemistry, Faculty of Sciences, University of Guilan, 41335-1914, Rasht, Guilan, Iran

^b Department of Chemistry, Faculty of Science, Tarbiat Modares University, Tehran, Iran

^c Young Researchers and Elites Club, Science and Research Branch, Islamic Azad University, Tehran, Iran

^d Department of Chemistry, Isfahan University of Technology, Isfahan, 84156-83111, Iran

* Corresponding author Tel.: +98-13-33367262; fax: +98-13-33367066 E-mail address: <u>khorshidi@guilan.ac.ir</u>

Keywords: Schiff base; Stöber method; carbon spheres; oxygen reduction reaction




Fig. S1. (a) Elemental mapping analysis of all, (b) C, (c) N, and (d) O elements of the SF-C resin spheres.




Fig. S2. (a) Elemental mapping analysis of all, (b) C, (c) N, and (d) O elements of the SF-P-C carbon spheres.