

Supplementary data

New Journal of Chemistry

Manuscript ID NJ-ART-05-2021-002663

***Cordyline fruticosa* (L.) A. Chev. Leaves: Isolation, HPLC/MS profiling and evaluation of nephroprotective and hepatoprotective activities supported by molecular docking.**

Mona Raslan^{1,*}, Rehab F. Taher², Ahmed A. Al-Karmalawy³, Dalia El-Ebeedy⁴, Angham G. Metwaly⁵, Nourhan M. Elkateeb⁶, Aml Ghanem⁷, Reem A. Elghaish⁸, Ahmed I. Abd El Maksoud⁹

¹ Pharmacognosy Department, National Research Centre, Dokki, 12622 Giza, Egypt.

² Chemistry of Natural Compounds Department, National Research Centre, 12622 Giza, Egypt.

³ Department of Pharmaceutical Medicinal Chemistry, Faculty of Pharmacy, Horus University-Egypt, New Damietta 34518, Egypt.

⁴ Pharmaceutical Biotechnology Department, Faculty of Biotechnology, Misr University for Science and Technology, Giza, Egypt.

⁵ Misr University for Science and Technology, Giza, Egypt.

⁶ Misr University for Science and Technology, Giza, Egypt.


⁷ Department of Molecular Biology, Genetic Engineering and Biotechnology Research Institute, University of Sadat City, Sadat City, Egypt.


⁸ Zewail city for science and technology, Giza, Egypt.



⁹ Industrial Biotechnology Department, Genetic Engineering and Biotechnology Research Institute, University of Sadat City, Sadat City, Egypt.

* Corresponding author. E-mail address: monaazzam78@gmail.com


Figure SI 1: 2 D pictures representing the binding interactions of the tested isolated compounds from *Cordyline fruticosa* L. (**1-12**) inside the binding pocket of the TNF- α protein compared to both silymarin (**13**) and the docked co-crystallized inhibitor (**14**).









8




9




10




11



12



13



14

