

Betaine Host-Guest Complexation with a Calixarene Receptor: Enhanced *In Vitro* Anticancer effect

Sherif Ashraf Fahmy¹ Fortuna Ponte² Iten M. Fawzy³ Emilia Sicilia², , Hassan Mohamed El-Said Azzazy^{1,*}

¹ Department of Chemistry, School of Sciences & Engineering, The American University in Cairo, AUC Avenue, PO. Box 74, New Cairo 11835, Egypt; sheriffahmy@aucegypt.edu

² Department of Chemistry and Chemical Technologies, University of Calabria, Arcavacata di Rende, 87036, Italy; fortuna.ponte@unical.it; emilia.sicilia@unical.it

³ Pharmaceutical Chemistry Department, Faculty of Pharmaceutical Sciences and Pharmaceutical Industries, Future University in Egypt, Cairo 12311, Egypt; iten.mamdouh@fue.edu.eg

* Correspondence: hazzazy@aucegypt.edu (H.M.E.-S.A.);

Tel.: +2 02 2615 2559; Fax: +2 02 2795 7565 (H.M.E.-S.A.)




Figure S1. ^1H NMR spectra of (A) individual *p*-SC4, (B) individual betaine, and (C) equimolar ratio of amount of betaine and *p*-SC4. Structures of *p*-SC4 and betaine are shown with appropriate protons labeled.




Figure S2. Absorbance spectra of several mixtures containing increasing concentrations of betaine (0.01–0.25 mM) and a fixed concentration of 0.2 mM *p*-SC4; all in distilled water.

