

Electronic Supplementary Material (ESI) for New Journal of Chemistry.

This journal is © The Royal Society of Chemistry and the Centre National de la Recherche Scientifique 2022

Supporting Information

Ternary photocatalysts based on MOF-derived TiO₂ co-decorated with ZnIn₂S₄ nanosheets and CdS nanoparticles for effective visible light degradation of organic pollutants

Huan Liu^{ab}, Jingzhe Li^{ab}, Yue Chen^{ab}, Xuting Sun^b, Xun Xu^{ab}, Lingfang Qiu^{ab}, Shuwang Duo^{*ab} and Ping Li^{*ab}

^aJiangxi Key Laboratory of Surface Engineering, Jiangxi Science and Technology Normal University,

Nanchang, Jiangxi 330013, P. R. China, dsw@jxstnu.edu.cn, lp1849065552@163.com

^bSchool of Materials and Mechanical & Electrical Engineering, Jiangxi Science and Technology Normal University, Nanchang 330013, P. R. China




Fig. S1 SEM image of MIL-125.




Fig. S2 EDS mapping of CdS/ZnIn₂S₄/TiO₂-II.




Fig. S3 XPS survey scan spectrum of CdS/ZnIn₂S₄/TiO₂-II.




Fig. S4 Recycling experiments of photocatalytic TC degradation.




Fig. S5 XRD patterns of CdS/ZnIn₂S₄/TiO₂-II before and after 5-cycle test in the photodegradation of RhB (a) and TC (b).




Fig. S6 SEM images of CdS/ZnIn₂S₄/TiO₂-II after 5-cycle test in the photodegradation of RhB (a) and TC (b).