

Supplementary Information

Structural and theoretical study of copper(II) complex incorporating chalcone and 2,2'-bipyridine mixed ligands : A probable candidate for optical material

Ruchi Gaur^{a#}, Parashuram Kallem^{b,c}*

^aDepartment of Chemistry, Institute of Science, Banaras Hindu University, Varanasi-221005
India

^bCenter for Membranes and Advanced Water Technology (CMAT), Khalifa University of
Science and Technology, P.O. Box 127788, Abu Dhabi, United Arab Emirates

^cDepartment of Chemical Engineering, Khalifa University of Science and Technology, PO
Box 127788, Abu Dhabi, United Arab Emirates

[#]Current address: Solid State and Structural Chemistry Unit, Indian Institute of Science,
Bangalore-560012, India

SI 1: Selected bond lengths (Å) and bond angles (°) of CuM.

Cu(1)-O(1)	1.880(2)	O(1)-Cu(1)-O(2)	92.23(9)
Cu(1)-O(2)	1.9193(19)	O(1)-Cu(1)-O(3)	99.95(19)
Cu(1)-O(3)	2.443(8)	N(1)-Cu(1)-N(2)	81.02(11)
Cu(1)-N(1)	1.992(3)	O(2)-Cu(1)-N(1)	92.55(10)
Cu(1)-N(2)	2.002(3)	O(3)-Cu(1)-N(1)	81.4(2)
S(1)-C(24)	1.703(5)	C(20)-S(1)-(24)	90.8(2)

SI 2: Geometric features of π - π interactions obtained from **CuM**.

Cg(I)… Cg(ring J)	Cg…Cg (Å)	α (°)	β (°)	γ (°)	Cg(I)…Prep (Å)	Cg(J)…Prep(Å)
Cg(4)-Cg(6) ^a	3.6783(19)	4.88	26.30	21.4	-3.4239(14)	-3.2975(11)

Symmetry transformation^a= -X,-Y,1-Z; Cg(4) = center of gravity of the ring N(1)-C(1)-C(2)-C(3)-C(4)-C(5); Cg(6) = center of gravity of the ring C(11)-C(12)-C(13)-C(14)- C(15)-C(16).