

Supporting Information

2D MXene interfaces preserve the basal electrophysiology of targeted neural circuits

Miao Xiao^{1, 3, 6, 7, #}, *Xiaoyun Li*^{1, 3, #}, *Simone Pifferi*³, *Beatrice Pastore*³, *Yun Liu*⁴, *Marco Lazzarino*⁸, *Vincent Torre*^{3, *}, *Xiaowei Yang*^{5, *}, *Anna Menini*^{3, *}, *Mingliang Tang*^{1, 2, *}

¹Institute for Cardiovascular Science & Department of Cardiovascular Surgery of the First Affiliated Hospital, Medical College, Soochow University, Suzhou, 215000, China.

²Co-innovation Center of Neuroregeneration, Nantong University, Nantong 226001, China

³Neurobiology Sector, International School for Advanced Studies (SISSA), via Bonomea 265, Trieste 34136, Italy.

⁴Key Laboratory for Developmental Genes and Human Disease, Ministry of Education, Institute of Life Sciences, Southeast University, Nanjing 210096, China

⁵School of Chemistry and Chemical Engineering, Frontiers Science Center for Transformative Molecules, Shanghai Jiao Tong University, Shanghai, China

⁶Suzhou Fishseeds Bio-technology, LTD, Suzhou 215138, China

⁷Anhui Isotex Biotech Co. Xuancheng 242300, China

⁸IOM-CNR, Area Science Park, Basovizza, Trieste 34149, Italy.


[#]These authors contributed equally to this work.

Email: torre@sissa.it, yangxw@sjtu.edu.cn, menini@sissa.it, mltang@suda.edu.cn


Supplementary Videos

Supplementary Video_Coverslip_001. Representative Fluo-4 AM fluorescence calcium imaging of 9 hippocampal neuron cultures on coverslip for 8 days *in vitro*.

Supplementary Video_MXene_001. Representative Fluo-4 AM fluorescence calcium imaging of 9 hippocampal neuron cultures on uncoated $Ti_3C_2T_x$ MXene film for 8 days *in vitro*.



Supplementary Figure 1. Two types of events from mPSCs which was recorded from a voltage-clamped hippocampal neuron (at holding potential of -50 mV) grown on the poly-L-ornithine coated coverslip control were identified. A slow decay component and a fast decay component were extracted from the mPSCs. These components differ in decay time (A, KS = 0.15, P = 0.001, Kolmogorov Smirnov test), amplitude (B, KS = 0.18, P = 0.009, Kolmogorov Smirnov test), and inter-event interval (C, KS = 0.55, P = 0.000). Slow (component 1) and fast (component 2) components, respectively referred to mIPSCs and mEPSCs.



Supplementary Figure 2. Representative SEM image of the Ti₃C₂T_x MXene film with enlarged higher-magnification. Scale bar = 5 μ m.