Electronic Supplementary Material (ESI) for RSC Advances. This journal is © The Royal Society of Chemistry 2022

Uncropped Western blot data

Structure-based design and synthesis of a novel long-chain 4"-alkyl ether derivative of EGCG as potent EGFR inhibitor: *In-vitro* and *in-silico* studies

Satyam Singh¹, Revathy Sahadevan², Rajarshi Roy¹, Mainak Biswas³, Priya Ghosh¹, Parimal Kar¹, Avinash Sonawane¹, Sushabhan Sadhukhan^{2,4,5*}

¹Department of Biosciences and Biomedical Engineering, Indian Institute of Technology Indore, Madhya Pradesh 453 552, India.

²Discipline of Chemistry, Indian Institute of Technology Palakkad, Kerala 678 623, India.

³School of Biotechnology, KIIT Deemed to be University, Bhubaneswar, Orissa 751 024, India.

⁴Physical & Chemical Biology Laboratory, Indian Institute of Technology Palakkad, Kerala 678 623, India.

⁵Department of Biological Sciences & Engineering, Indian Institute of Technology Palakkad, Kerala 678 623, India.

*Corresponding author. E-mail address: sushabhan@iitpkd.ac.in

Fig. 3.





Fig. 6.










Fig. 7.













EGFR







ERK Gapdh











β-actin





pAkt β-actin