

Electronic Supplementary Information

Orientation Control of Lyotropic Chromonic Liquid Crystals in the Capillary Bridge

Hee Seong Yun,^a Geonhyeong Park^a and Dong Ki Yoon^{*ab}

a Department of Chemistry, Korea Advanced Institute of Science and Technology (KAIST), Daejeon, 34141, Republic of Korea

b Graduate School of Nanoscience and Technology and KAIST Institute for Nanocentury, Korea Advanced Institute of Science and Technology (KAIST), Daejeon, 34141, Republic of Korea

Legends

Fig. S1 Synthetic route of PTCDA to PDI-HCl

Fig. S2 Schematic illustrations of the formation of the capillary bridges in the sandwich cell

Fig. S3 POM images of uniaxially aligned PDI-HCl in the sandwich cell with diamond-scratched silicon and glass

Movie S1. Phase transition to N phase of PDI-HCl in the capillary bridge at $w = 20 \mu\text{m}$

Movie S2. Phase transition to N phase of PDI-HCl in the capillary bridge at $w = 50 \mu\text{m}$

Movie S3. Phase transition to N phase of PDI-HCl in the capillary bridge at $w = 100 \mu\text{m}$




Fig. S1 Synthetic route of PTCDA to PDI-HCl




Figure S2. Schematic illustrations of the formation of the capillary bridges in the sandwich cell




Figure S3. (a,b,c) POM and (d,e,f) POM with λ plate images of uniaxially aligned PDI-HCl in the sandwich cell with diamond-scratched silicon and glass. Red arrow means scratch direction and PDI-HCl aggregates aligned parallel to scratch direction.¹⁹ Scale bar is 100 μm .