

1 **Supporting Information**

2 **Structure relationship of non-covalent interactions between lotus seedpod**

3 **oligomeric procyanidins and glycated casein hydrolysate during digestion**

4 Yu Ouyang^{a,b}, Yinggang Liang^a, Mengyao Niu^a, Jia Yan^a, Qianmei Chu^c, Mengzhou

5 Zhou^a, Wei Li^a, Nianjie Feng^{a,*}, Qian Wu^{a,*}

6 a Cooperative Innovation Center of Industrial Fermentation (Ministry of Education &
7 Hubei Province), Key Laboratory of Fermentation Engineering (Ministry of
8 Education), National “111” Center for Cellular Regulation and Molecular
9 Pharmaceutics, Hubei Key Laboratory of Industrial Microbiology, Hubei University
10 of Technology, Wuhan, Hubei, 430068, China

11 b Agricultural College, Hubei Three Gorges Polytechnic, Yichang 443000, PR China

12 c Xiangyang Academy of Agricultural Sciences, Xiangyang, Hubei 441057, PR
13 China


14 Email addresses: Yu Ouyang (oyy940098506@163.com), Yinggang Liang
15 (liangyinggang2021@163.com), Mengyao Niu (nmy18239532535@163.com), Jia
16 Yan (yj199721@163.com), Qianmei Chu (phya49@163.com), Mengzhou Zhou
17 (zmzkelvin@163.com), Wei Li (ucee71@163.com)

18 Corresponding author: Email: Qian Wu (wuqian@hbut.edu.cn), Nianjie Feng
19 (njfeng@hbut.edu.cn), Tel numbers: +86-13720167990

20 Mail address: Hubei University of Technology, Wuhan, Hubei, 430068, China

Contents:

Supporting Information Figure S1. The curve-fitting spectra of amide I band (1700-1600 cm^{-1}) of the GCSH-procyanidin complex.



Supporting Information Figure S1