

Insights into the catalytic properties of 4,3- α -glucanotransferase to guide the biofabrication of α -glucans with low digestibility

Yuqi Yang ¹, Yujing Sun ¹, Tao Zhang ¹, Bruce R. Hamaker ^{1,2}, Ming Miao ^{1*}

¹ *State Key Laboratory of Food Science & Technology, Jiangnan University, 1800 Lihu Avenue, Wuxi, Jiangsu 214122, P. R. China*

² *Whistler Center for Carbohydrate Research, Purdue University, 745 Agriculture Mall Drive, West Lafayette, IN 47907, United States*

* Corresponding author. Address: State Key Laboratory of Food Science & Technology, Jiangnan University, 1800 Lihu Avenue, Wuxi, Jiangsu 214122, P. R. China.

Tel : +86 (0)510 853 27859, Fax: +86 (0)510 859 19161.

E-mail address: miaoming@jiangnan.edu.cn (Miao M.)





Fig.S1. HPAEC profile of MD-2, MD-6, and MD-29.

(a)

Apr04-2023
MM-Y1Q-001



(b)

Apr04-2023
MM-Y1Q-005




Fig.S2. ¹H NMR spectra of MD-2 and 4,3- α -GTase treated MD-2 for 72 h.