Electronic Supplementary Material (ESI) for RSC Medicinal Chemistry. This journal is © The Royal Society of Chemistry 2024

Inhibition of DXR in the MEP Pathway with Lipophilic N-alkoxyaryl FR900098 Analogs


Darean Bague^{†,a}, Ruiqin Wang^{†,a}, Dana Hodge[§], Marwa O. Mikati[#], Jose S. Roma[‡], Helena I. Boshoff[‡], Allyson L. Dailey[⊥], Misgina Girma[⊥], Robin D. Couch[⊥], Audrey R. Odom John^{#,§}, and Cynthia S. Dowd*,[†]

Supplementary Information


Figure S1. IC ₅₀ curves for <i>P. falciparum</i> Dxr inhibition	S2
Figure S2. IC ₅₀ curves for <i>M. tuberculosis</i> Dxr inhibition	S6
Figure S3. IC ₅₀ curves for wild-type and <i>had1</i> mutant <i>P. falciparum</i> inhibition	S7
Figure S4. Comparison charts for wild-type and <i>had1</i> mutant <i>P. falciparum</i> inhibition	S9

Figure S1. IC₅₀ curves for *P. falciparum* Dxr inhibition


Compound 6a




Compound 6e




Compound 6f




Compound 6j



Compound 6k



Compound 6I



Compound 6n





Figure S2. IC_{50} curves for *M. tuberculosis* Dxr inhibition

Compound 6I



Compound 6n

