

SUPPORTING INFORMATION

For

**Identification of Novel Phenylalanine Derivatives Bearing Hydroxamic
Acid moiety as Potent Quorum Sensing Inhibitors.**

Truong Thanh Tung,^{1,2,*} Nguyen Quoc Thang,^{3,4} Nguyen Quang Huy,¹ Pham Bao
Phuong,¹ Dinh Ngoc Minh,¹ Nguyen Hai Nam,³ John Nielsen⁵

¹*Faculty of Pharmacy, PHENIKAA University, Hanoi 12116, Vietnam*

²*PHENIKAA Institute for Advanced Study (PIAS), PHENIKAA University, Hanoi 12116, Vietnam*

³*Hanoi University of Pharmacy, 13-15 Le Thanh Tong, Hanoi, Vietnam*

⁴*Vinmec International Hospital, Hanoi, Vietnam*

⁵*Department of Drug Design and Pharmacology, University of Copenhagen, Denmark*



*Corresponding author, Email: tung.truongthanh@phenikaa-uni.edu.vn

ORCID: 0000-0002-5263-203X



tunglab.com


COPY OF NMR SPECTRA



Key intermediate 2:







Carboxylic acid intermediates:
































Compound 4a











Compound **4b**








Compound 4c








Compound **4d**








Compound 4e








Compound 4f








Compound 4g





Compound 4h





Compound 4i

