

2-aminofluorene based triazole cocktailed with organosilane as UV-Visible and fluorescence sensor for highly selective detection of Fe(III) ion and its anti-leishmanial activity

Gurjaspreet Singh^{a}, Sofia Gupta^{a*}, Pawan^a, Mohit^a, Diksha^a, Isha Saini^b, Vikas^a, Brij Mohan^c, Sanjeev Soni^d*

^aDepartment of Chemistry, Panjab University, Chandigarh, 160014, India

^bDepartment of Zoology, Panjab University, Chandigarh, 160014, India

^cCollege of Ocean Food and Biological Engineering, Jimei University, 185 Yinjiang Road, Jimei District, Xiamen, 361021

^dDepartment of Chemistry, GGSDS college Chandigarh, 160032 India

*Corresponding Authors

Prof. Gurjaspreet Singh

Department of Chemistry & Centre of Advanced Studies

Panjab University, Chandigarh, India

Email: gjpsingh@pu.ac.in

Tel. No: 0172-2534428, 09814302099, 9317502099

Fax No: 0172-2545074

Miss Sofia Gupta

Research Scholar

Department of Chemistry & Centre of Advanced Studies

Panjab University, Chandigarh, India

Email: sofiagupta02@gmail.com

Tel. No: 9888120902

Table of Contents :

Entry	Page No.
¹ H NMR spectrum of compound 5a (Fig.S1).....	3
¹ H NMR spectrum of compound 5b (Fig.S2).....	3
¹ H NMR spectrum of compound 6a (Fig.S3).....	4
¹ H NMR spectrum of compound 6b (Fig. S4).....	4
¹³ C NMR spectrum of compound 5a (Fig. S5).....	5
¹³ C NMR spectrum of compound 5b (Fig. S6).....	5
¹³ C NMR spectrum of compound 6a (Fig. S7).....	6
¹³ C NMR spectrum of compound 6b (Fig. S8).....	6
Mass spectrum of compound 5a (Fig. S9).....	7
Mass spectrum of compound 5b (Fig. S10)	7
Mass spectrum of compound 6a (Fig.S11).....	8
Mass spectrum of compound 6b (Fig.S12).....	8
pH effect of 6a and 6a+Fe ³⁺ complex (Fig.S13).....	9
Selectivity studies (Fig. S14).....	9
Response time studies (Fig. S15).....	9
¹ H NMR spectra of compound 6a and 6a+Fe ³⁺ (Fig.S16).....	10
XPS spectra of 6a and 6a+Fe ³⁺ of C 1s and Si 2p (Fig.S17).....	10




Fig.S3: ^1H NMR spectrum of compound **6a**




Fig.S4: ^1H NMR spectrum of compound **6b**




Fig.S5: ^{13}C NMR spectrum of compound **5a**




Fig.S6: ^{13}C NMR spectrum of compound **5b**




Fig.S7: ^{13}C NMR spectrum of compound **6a**






Fig.S8: ^{13}C NMR spectrum of compound **6b**

**Fig.S9: Mass spectrum spectrum of compound 5a****Fig.S10: Mass spectrum spectrum of compound 5b**

WATERS,Q-TOF MICROMASS (ESI-MS)
SOFIA SIL_S_1 17 (0.232) Sm (SG, 2x3.00); Cm (2:28)

SAIF/CIL,PANJAB UNIVERSITY,CHANDIGARH

TOF MS ES+
1.51e4




Fig.S11: Mass spectrum spectrum of compound 6a

SAIF,PANJAB UNIVERSITY CHANDIGARH

SYNAPT-XS#DBA064

20-Jun-2022
15:20:09
1: TOF MS ES+
4.45e4




Fig.S12: Mass spectrum spectrum of compound **6b**




Fig.S13: pH effect




Fig.S14: Selectivity studies




Fig.S15: Response time studies




Fig.S16 ^1H NMR spectra of compound **6a** and its complex with Fe^{3+} ions.




Fig.S17 XPS spectra of **6a** and 6a+Fe³⁺. (a) C 1s spectra of **6a**, (b) C 1s spectra of 6a+Fe³⁺, (c) Si 2p spectra of **6a**, and (d) Si 2p spectra of 6a+Fe³⁺.