

Supporting Information

External Electric field Driven Electronic Structures: Tunable Nonlinear Optical Properties in Helical Bilayer Nanographenes

Ping-yao Gan,^a Xiao Huang,^a Feng-wei Gao,^{*ab} Hong-liang Xu^{*c}

^a School of Chemistry and Environmental Engineering, Changchun University of Science and Technology, 7989 Weixing Road, Changchun 130012, China.

^b Chongqing Research Institute, Changchun University of Science and Technology, No.618 Liangjiang Avenue, Longxing Town, Yubei District, Chongqing City 401135, China.

E-mail: gaofw@cust.edu.cn

^c Institute of Functional Material Chemistry, Department of Chemistry, Northeast Normal University, Changchun 130024, China.

E-mail: hlxu@nenu.edu.cn

Table of contents

Figure S1. Evolutions of the non-covalent interaction (NCI) and reduced density gradient (RDG) maps of the [9]HBNG under the vertical F_z ($F_z = 1 \times 10^{-4}$, au)	2
Figure S2. Evolutions of dipole moment (μ) of ground state under the vertical F_z ($F_z = 1 \times 10^{-4}$, au)	3
Figure S3. Evolutions of the first hyperpolarizabilities (β_{tot}) of [9]HBNG under the vertical F_z ($F_z = 1 \times 10^{-4}$, au) at the M06-2X/6-31G**, CAM-B3LYP/6-31G** and BHandHLYP/6-31G** levels	4
Figure S4. Evolutions of the electron density difference maps (EDDM) of the [9]HBNG under the vertical F_z ($F_z = 1 \times 10^{-4}$, au)	5
Figure S5. Evolutions of the 2D second-order NLO spectra the [9]HBNG under the F_z ($F_z = 1 \times 10^{-4}$, au)	6
Figure S6. Evolutions of the electron circular dichroism (ECD) spectrum of the [9]HBNG under the F_z ($F_z = 1 \times 10^{-4}$, au)	7
Table S1. Evolutions of the β_x , β_y , β_z and β_{tot} values of [9]HBNG under the vertical F_z ($F_z = 1 \times 10^{-4}$, au)	8

Supporting Information




Figure S1. Evolutions of the non-covalent interaction (NCI) and reduced density gradient (RDG) maps of the [9]HBNG under the vertical F_z ($F_z = 1 \times 10^{-4}$, au).




Figure S2. Evolutions of dipole moment (μ) of ground state under the vertical F_z ($F_z = 1 \times 10^{-4}$, au).




Figure S3. Evolutions of the first hyperpolarizabilities (β_{tot}) of [9]HBNG under the vertical F_z ($F_z = 1 \times 10^{-4}$, au) at the M06-2X/6-31G**, CAM-B3LYP/6-31G** and BHandHLYP/6-31G** levels.




Figure S4. Evolutions of the electron density difference maps (EDDM) of the [9]HBNG under the vertical F_z ($F_z = 1 \times 10^{-4}$, au).




Figure S5. Evolutions of the 2D second-order NLO spectra the [9]HBNG ((a) $F_z = 0$ au, (b) $F_z = 100 \times 10^{-4}$ au).




Figure S6. Evolutions of the electron circular dichroism (ECD) spectrum of the [9]HBNG under the F_z ($F_z = 1 \times 10^{-4}$, au).

Supporting Information

Table S1. Evolutions of the β_x , β_y , β_z and β_{tot} values of [9]HBNG under the vertical F_z ($F_z = 1 \times 10^{-4}$, au).

$F_z = 1 \times 10^{-4}$ (au)	β_x	β_y	β_z	β_{tot}
0	653	0	0	653
20	654	143	538	859
40	670	337	1121	1349
60	689	584	2028	2219
80	707	1011	4369	4540
100	1023	1603	12434	12579