

SUPPORTING INFORMATION

Modification of Hydrothermally Synthesized α - Fe_2O_3 Nanorods with $\text{g-C}_3\text{N}_4$ Prepared from Various Precursors as Photoanodes for Hydrogen Production

Muhammad Ibadurrohman^{*1)}, Afaf Qurrotu Ainin¹⁾, Fakhri Zinul Alam¹⁾, Nadia Mumtazah¹⁾, Slamet¹⁾, Alfian Ferdiansyah Madsuha²⁾, Reza Miftahul Ulum²⁾, and Bonavian Hasiholan³⁾

1) Department of Chemical Engineering, Faculty of Engineering, Universitas Indonesia, Depok, Indonesia 16424

2) Department of Metallurgical and Material Engineering, Faculty of Engineering, Universitas Indonesia, Depok, Indonesia 16424

3) Department of Chemical & Petroleum Engineering, UCSI University, 56000 Kuala Lumpur, Malaysia

*corresponding author: ibad@che.ui.ac.id




Figure S1. Longer operation of light-chopped chronoamperometric measurements α - $\text{Fe}_2\text{O}_3/\text{g-C}_3\text{N}_4$ performed in three cycles (reused) without glycerol as a sacrificial agent. Note: the normalized photocurrent is the photocurrents divided by the highest photocurrent data point in all three cycles.