

Electronic Supplementary Information (ESI)

Cyclic-hanging-multicyclic polymer synthesized by ladderphane-mediated blocking-cyclization technique for direct visualization of cyclic macromolecular topology


Cuihong Ma,^{a,b}† Ying Quan,^a† Xiaojuan Liao,^b Ruyi Sun^{b*} and Meiran Xie^{b*}

^aSchool of Chemistry and Molecular Engineering, East China Normal University, Shanghai 200241, China. E-mail: rysun@chem.ecnu.edu.cn, mrxie@chem.ecnu.edu.cn

^bMinistry of Education Key Laboratory of Macromolecular Synthesis and Functionalization, Department of Polymer Science and Engineering, Zhejiang University, Hangzhou, Zhejiang 310058, China.

†These authors contributed equally.

Supplementary Data



Scheme S1 Synthetic routes of **PMT** (a) and cyclic side chain-contained macromonomer **NB(*c*-PPMT)** (b).




Fig. S1 ^1H NMR (500 MHz) spectrum of **PMT** measured in CDCl_3 at 25 °C.




Fig. S2 ^1H NMR (500 MHz) spectrum of N-(4-hydroxymethylbiphenyl)-norbornene dicarboximide measured in CDCl_3 at 25 °C.




Fig. S3 ^1H NMR (a) and ^{13}C NMR (b) spectra of RNB measured in CDCl_3 at 25 °C.




Fig. S4 ¹H NMR (a) and ¹³C NMR (b) spectra of cyclic side chain-contained macromonomer **NB(c-PPMT)** measured in CDCl₃ at 25 °C.




Fig. S5 ¹H NMR spectrum of *c*-{PBNP₁₀-[*b*-P(TNP₁₉₅-*co*-NB(*c*-PPMT))₅]}₂-*b*-PBNP₁₀} in CDCl₃.




Fig. S6 ¹H NMR spectrum of PBNP₂₀ in CDCl₃.




Fig. S7 ¹H NMR spectrum of *l*-{PBNP₂₀-*b*-[P(TNP₁₉₅-*co*-NB(*c*-PPMT)₅)]₂} in CDCl₃.




Fig. S8 UV-vis spectra of polymers.




Fig. S9 DLS diagrams of cyclic polymers in THF at 2 mg mL⁻¹.