

Supporting Information

Terbium-doped carbon dots (Tb-CDs) as a novel contrast agent for efficient X-ray attenuation

Anara Molkenova^{1#}, Lazzat Serik^{2#}, Alina Ramazanova³, Kamila Zhumanova², Bakyt Duisenbayeva⁴, Ainur Zhussupbekova⁵, Kuanysh Zhussupbekov⁵, Igor V. Shvets⁵, Ki Su Kim^{1,7,8}, Dong-Wook Han⁶, Timur Sh. Atabaev^{2*}

¹*Institute of Advanced Organic Materials, Pusan National University, Busan 46241, Republic of Korea*

²*Department of Chemistry, Nazarbayev University, Astana 010000, Kazakhstan*

³*Department of Biology, University of Padua, Padua 35121, Italy*

⁴*Department of Radiology, Republican Diagnostic Center, Astana 010000, Kazakhstan*

⁵*School of Physics and Centre for Research on Adaptive Nanostructures and Nanodevices (CRANN), Trinity College Dublin, Dublin, Ireland*

⁶*Department of Cogno-Mechatronics Engineering, Pusan National University, Busan 46241, Republic of Korea*

⁷*School of Chemical Engineering, College of Engineering, Pusan National University, 2 Busandaehak-ro 63 beon-gil, Geumjeong-gu, Busan 46241, Republic of Korea*

⁸*Department of Organic Material Science & Engineering, Pusan National University, 2 Busandaehak-ro 63 beon-gil, Geumjeong-gu, Busan 46241, Republic of Korea*

- these authors contributed equally

*Correspondence: timur.atabaev@nu.edu.kz (T.S. Atabaev)




Figure S1. A) Phase contrast image of L929 cells incubated with Tb-CDs, and B) fluorescence image of the same region of interest




Figure S2. Morphological alterations of L929 cells incubated with Tb-CDs at different concentrations