

Supporting file for Injectable hydrogels of recombinant human collagen type III and chitosan with antibacterial and antioxidative activities for wound healing

Li Xiong^{a,b, &}, Chen Zhou^{a,b, &}, Lei Tong^{a,b}, Xiaowen Han^{a,b}, Yaping Zou^{a,b}, Zuqin Dong^{a,b}, Jie Liang^{*a,b,c}, Yafang Chen^{*a,b}, Yujiang Fan^{a,b}

^a National Engineering Research Center for Biomaterials, Sichuan University, 29 Wangjiang Road, Chengdu, Sichuan, 610064, PR China

^b College of Biomedical Engineering, Sichuan University, 29 Wangjiang Road, Chengdu, Sichuan, 610064, PR China

^c Sichuan Testing Center for Biomaterials and Medical Devices, Sichuan University, 29 Wangjiang Road, Chengdu, 610064, PR China

&. These authors contributed equally to this work.

* Corresponding authors at: Biomaterials Building, Sichuan University, Chengdu, 610064, China.

E-mail addresses: jie_L88@126.com, yafang.chen@scu.edu.cn

Results and discussion




Fig. S1. Statistics of pore diameter of hydrogels.




Fig. S2. The storage modulus of hydrogels.




Fig. S3. The survival rates of hydrogels against *E. coli* and *S. aureus*.




Fig. S4. H&E staining in different treatment groups of the major organs tissues on day 6.