

Supporting Information

Biocompatible Recombinant Type III Collagen Enhancing Skin Repair and Anti-Wrinkle Effects

Mingzhu Ye ^{a, c, d}, Yirui Fan ^{a, d}, Caihong Fu ^{a, d}, Huixia He, ^{b, d*} Jianxi Xiao ^{a, c, d*}

^aState Key Laboratory of Applied Organic Chemistry, College of Chemistry and Chemical Engineering, Lanzhou University, Lanzhou 730000, P. R. China

^bCollege of School of Life Sciences, Lanzhou University, Lanzhou 730000, P. R. China

^cCollege of Earth and Environmental Sciences, Lanzhou University, Lanzhou 730000, P. R. China

^dGansu Engineering Research Center of Medical Collagen, Lanzhou 730000, PR China

Email: xiaojx@lzu.edu.cn; hehuix@lzu.edu.cn.




Fig. S1 SEM image of the rCol III.




Fig. S2 Rheological properties of rCol III.




Fig. S3 The content of SOD(A) and MDA(B) of mice skin injection of different groups, including blank, vehicle, HA, rCol III-0.1, rCol III-0.1, and rCol III-1.0 groups.




Fig. S4 Quantifications of collagen content.




Fig. S5 Typical plot of zebrafish anti-wrinkle efficacy of blank group. Scale bar: 200 nm.

Table S1. Quantitative calculation of the inhibition rate of caudal fin shrinkage in zebrafish.

Groups	Caudal fin		p-Value	Q(%)
	area (pixel ²)	Mean ± SE		
Blank	1231264 ± 39074		<0.0001	-
Vehicle	458947 ± 68338		-	-
Polyphenols	774634 ± 90482		0.0141	40.88
rCol III-0.1 mg/mL	681055 ± 78462		0.0498	28.76
rCol III-0.5 mg/mL	857650 ± 97988		0.0046	51.62