

Supplementary Information

Electrochemical Kinetic Fingerprinting of Single-Molecule Coordinations in the Confined Nanopores

Chao-Nan Yang^[a], Wei Liu^[a], Hao-Tian Liu^[a], Ji-Chang Zhang^[a], Yi-Tao Long^[a], Yi-Lun Ying^{[a,b]*}

^a Molecular Sensing and Imaging Center, School of Chemistry and Chemical Engineering, Nanjing University, Nanjing 210023, P. R. China

^b Chemistry and Biomedicine Innovation Center, Nanjing University, Nanjing 210023, P. R. China

* E-mail: yilunying@nju.edu.cn

Table of Contents

Fig. S1. Current-voltage (<i>I-V</i>) curves of K238H-AeL, S276H-AeL and WT-AeL nanopore.....	3
Fig. S2. Control experiment of WT-AeL nanopore.....	4
Fig. S3. MD simulations of K238H-AeL nanopore and S276H-AeL nanopore.....	5
Table. S1. Distance of N atom to O atom in K238H-AeL or S276H-AeL.....	5
Author Contributions.....	6




Fig. S1. Current-voltage (*I-V*) curves of (a) K238H-AeL, (b) S276H-AeL and (c) WT-AeL aerolysin nanopore. The data were acquired in the condition of 1.0 M KCl, 10 mM Tris, and 1.0 mM EDTA at pH 8.0. The error bars are the standard error of the mean (s.e.m) for 3 independent experiments.




Fig. S2. Control experiment of WT-AeL nanopore (a) The current trace of WT-AeL nanopore treated with the 0.5 mM H₂AuCl₄, the current trace was acquired in the condition of 1.0 M KCl, 10 mM Tris, and 1.0 mM EDTA at pH 8.0. The applied voltage is 100 mV. No reaction occurs between WT-AeL nanopore and the H₂AuCl₄ after recording with 30 min; (b) The mass spectrometry of WT-AeL proaerolysin; (c) The mass spectrometry of WT-AeL proaerolysin with Au(III). The peaks at $m/z = 53387 \pm 1$ and $m/z = 53388 \pm 1$ are both assigned to unreacted proaerolysin WT-AeL proaerolysin [M]⁺. The concentration of Au(III) is 0.5 mM.




Fig. S3. MD simulations of K238H-AeL nanopore and S276H-AeL nanopore. (a) The top view of K238H-AeL nanopore. H238 is shown in blue, E258 site is shown in red. (b) The top view of S276H-AeL nanopore. H238 is shown in blue, D222 site is shown in red. The equilibrated time is 5 ns.

Table. S1. Distance of N atom to O atom in K238H-AeL or S276H-AeL

Distance of the N atom to O atom	$d(\text{K238H-AeL}) / \text{Å}^*$	$d(\text{S276H-AeL}) / \text{Å}^{\&}$
d_1	4.4	6.1
d_2	4.3	4.9
d_3	4.4	6.6
d_4	5.3	4.8
d_5	4.9	5.0
d_6	4.6	7.3
d_7	4.4	7.1
Average d^{\S}	4.6 ± 0.3	6.0 ± 1.0

* The distance of N atom of the imidazole ring in H238 site to O atom of carboxyl in E258 site under K238H-AeL nanopore.

& The distance of N atom of the imidazole ring in H276 site to O atom of carboxyl in D222 site under K276H-AeL nanopore.

\S The average $d = (d_1 + d_2 + d_3 + d_4 + d_5 + d_6 + d_7) / 7$

Author Contributions

Y.-L. Ying and Y.-T. Long conceived idea. C.-N. Yang and conducted the experiment. C.-N. Yang analyzed the data. C.-N. Yang, W. Liu, H.-T. Liu and J.-C. Zhang prepared the manuscript. Y.-L. Ying and Y.-T. Long wrote the manuscript.