Supplementary Data

Dietary supplementation of pterostilbene, the major component from small berries, prevents alcohol induced liver injury associated with lipid accumulation and inflammation

Jia-Yi Dou^{a, 1}, Sai-Hu Liu^{a, 1}, Jia Guo^a, Chen-Yu Wang^a, Xu Dai^a, Li-Hua Lian^a,

Zhen-Yu Cui^{a,b,*}, Ji-Xing Nan^{a,*}, Yan-Ling Wu^{a,*}

^a Key Laboratory for Traditional Chinese Korean Medicine Research (State Ethnic Affairs), College of Pharmacy, Yanbian University, Yanji, Jilin Province 133002, China

^b Jilin Vocational and Technical College, Yanji, Jilin Province 133002, China

*Corresponding authors at: Key Laboratory for Traditional Chinese Korean Medicine Research (State Ethnic Affairs), College of Pharmacy, Yanbian University, Yanji, Jilin Province 133002, China

E-mail addresses: <u>cuizhenyu@ybu.edu.cn</u> (Z.-Y. Cui); <u>jxnan@ybu.edu.cn</u> (J.-X. Nan); <u>ylwu@ybu.edu.cn</u> (Y.-L. Wu).




Fig. S1 (A) Chemical structure of PTE. (B) Animal experimental scheme of acute ethanol model. (C) Animal experimental scheme of chronic ethanol-binge feeding model.

Normal group: control diet

pair-fed group: control liquid diet

(change diets at 3:00-5:00 pm)

9h

Isocaloric

maltose dextrin gavage

Euthanize

8d control liquid diet

ad libitum