

Supplementary Material (ESI) for Lab on a Chip

Deciphering Hepatoma Cell Resistance to Tyrosine Kinase Inhibitors: Insights from a Liver-on-a-Chip Model Unveiling Tumor Endothelial Cell Mechanisms

Madhu Shree Poddar,^a Yu-De Chu,^{bcd} Chau-Ting Yeh,^{bcd} Cheng-Hsien Liu^{*aef}

^aInstitute of Nanoengineering and Microsystems, National Tsing Hua University, Hsinchu, 30044, Taiwan, R.O.C.

^bLiver Research Center, Chang Gung Memorial Hospital, Taoyuan 333, Taiwan, R.O.C.

^cMolecular Medicine Research Center, Chang Gung University, Taoyuan 333, Taiwan, R.O.C.

^dInstitute of Stem Cell and Translational Cancer Research, Chang Gung Memorial Hospital, Taoyuan 333, Taiwan, R.O.C


^eDepartment of Power Mechanical Engineering, National Tsing Hua University, Hsinchu 30044, Taiwan, R.O.C.

^fCollege of Semiconductor Research, National Tsing Hua University, Hsinchu 30044, Taiwan, R.O.C.

*E-mail: liuch@pme.nthu.edu.tw


Supplementary Figures

Figure S1.



The IC₅₀ values of Lenvatinib and Sorafenib are 20.7μM and 10.4 μM, respectively.

Figure S2.



The step by step working process for our liver-chip.