

Electronic Supplementary Information

Acoustic enrichment of sperm for in vitro fertilization

Chunqiu Zhang,^{a,c,†} Ning Rong,^{a,†} Ziyi Lin,^{b,c,†} Peng-Qi Li,^a Jingyao Shi,^a Wei Zhou,^a
Lili Niu,^a Fei Li,^a Rongxin Tang,^{*e} Lei Li^{*b} and Long Meng^{*a,d}

^aKey Laboratory of Biomedical Imaging Science and System, Shenzhen Institutes of Advanced Technology, Chinese Academy of Sciences, Shenzhen, Guangdong 518055, P. R. China.

^bThe Brain Cognition and Brain Disease Institute, Shenzhen Institute of Advanced Technology, Chinese Academy of Sciences, Shenzhen, P. R. China.

^cUniversity of Chinese Academy of Sciences, Beijing 100190, P. R. China.

^dCollege of Medicine and Biological Information Engineering, Northeastern University, Shenyang, Liaoning 110016, P. R. China

^eCenter for Reproductive Medicine, Shanghai 10th People's Hospital of Tongji University, Shanghai 200072, P. R. China.

[†]These authors contributed equally to this work.

E-mail: long.meng@siat.ac.cn; Tel: +86 755 86392484

Supplementary Figures.




Fig. S1 The relative position of the liquid droplet in the device.




Fig. S2 Enrichment process of live and dead sperm around PS particles with a diameter of $100 \mu\text{m}$ during acoustic treatment. (a) Fluorescently stained sperm with calcein-AM, which indicated their viability. (b) Fluorescently stained sperm with PI, which indicates their death. (c) The merged images of fluorescently stained sperm with calcein-AM and PI.




Fig. S3 The temperature distribution of a 1 μL droplet on the substrate over time.