

Selective Induction of Apoptotic Cell Death in Lung Carcinoma Cells by Curcumin loaded PEGylated Lipid Nanoparticles with Minimal Normal Tissue Toxicity: *In-Vitro* & *In-Vivo* Toxicity Evaluation by Oral Delivery

Bijaideep Dutta^{a,b}, Sourav Kumar Das^{b,c}, Mayur Temgire^{d,e}, Jayesh Bellare^e, K. C. Barick^{a,b*}, Amit Kumar^{b,c}, P. A. Hassan^{a,b*}

^aChemistry Division, Bhabha Atomic Research Centre, Trombay, Mumbai 400085, India

^bHomi Bhabha National Institute, Anushaktinagar, Mumbai 400094, India

^cRadiation Biology and Health Science Division, Bhabha Atomic Research Centre, Trombay, Mumbai 400085, India

^dAdvanced Centre for Treatment, Research and Education in Cancer (ACTREC), Navi Mumbai-410210, India

^e Indian Institute of Technology, Bombay, Powai, Mumbai-400076, India

*Corresponding author: Tel.: + 91 22 2559 0284, Fax: + 91 22 2550 5099,
Email: kbarick@barc.gov.in (K. C. Barick), hassan@barc.gov.in (P. A. Hassan)

Supplementary data




Fig. 1S: Percentage proliferation of A549 cells after 24h (A) or 48h (B) of treatment with CSLN (1-50 μM) as determined by MTT assay. The values are mean±SD for three independent experiments performed in triplicates. **p<0.01, ***p<0.001, ****p<0.0001 compared to control group. ns: Not significant.