

SUPPORTING INFORMATION

Redox-Responsive Micellar Nanoparticles Using Benzothiazole-disulfide Terminated Polymers: Employing Host-Guest Complexation for Targeted Delivery of Curcumin

*Salli Kocak,^a Beyza Demirkol,^a Ruveyda Kilic Boz,^a Rana Sanyal,^{*a,b}*

*Mehmet Arslan,^{*c} Amitav Sanyal^{*a,b}*

^a Department of Chemistry, Bogazici University, Bebek, Istanbul, 34342, Türkiye

^b Center for Life Sciences and Technologies, Bogazici University, Istanbul, Türkiye

^c Department of Polymer Materials Engineering, Faculty of Engineering, Yalova University, Yalova 77200, Türkiye

List of Figures

Fig. S1 ¹³C NMR spectrum of the BDS-bearing CTA.

Fig. S2 SEC traces of P1, P2, and P3 polymers.

Fig. S3 SEC traces of P1 and CD-PEG polymers.

Fig. S4 ¹H NMR spectrum of the polymer CD-PEG-COOH in CDCl₃.

Fig. S5 SEC traces of CD-PEG, CD-PEG-COOH and CD-PEG-RGD polymers.

Fig. S6 The size distribution histograms of curcumin-loaded nanoparticles according to STEM image.

Fig. S7 UV spectra of curcumin released from nanoparticles in 5 mM GSH in Acetate buffer (pH 5.5) and in 5 mM GSH in PBS buffer (pH 7.4)

Fig. S8 (a) DLS analysis of curcumin-loaded nanoparticles in PBS; (b) after 4 hours, and (c) after 24 hours in GSH solution (0.010 M).

Fig. S9 (a) Flow cytometry data and (b) fluorescence microscopy images of U-87 MG cell line treated with curcumin encapsulated targeted nanoparticles for 4 hours at 4 °C and 37 °C. Scale bar: 20 μm.




Fig. S1 ^{13}C NMR spectrum of the BDS-bearing CTA.




Fig. S2 SEC traces of P1, P2, and P3 polymers.




Fig. S3 SEC traces of P1 and CD-PEG polymers.




Fig. S4 ¹H NMR spectrum of the polymer CD-PEG-COOH in CDCl₃.




Fig. S5 SEC traces of CD-PEG, CD-PEG-COOH and CD-PEG-RGD polymers.




Fig. S6 The size distribution histograms of curcumin-loaded nanoparticles according to STEM image.




Fig. S7 UV spectra of curcumin released from nanoparticles in 5 mM GSH in Acetate buffer (pH 5.5) and in 5 mM GSH in PBS buffer (pH 7.4).




Fig. S8 (a) DLS analysis of curcumin-loaded nanoparticles in PBS; (b) after 4 hours, and (c) after 24 hours in GSH solution (0.010 M).




Fig. S9 (a) Flow cytometry data and (b) fluorescence microscopy images of U-87 MG cell line treated with curcumin encapsulated targeted nanoparticles for 4 hours at 4 °C and 37 °C.