Synthesis and Bio-evaluation of Aminoferrocene-Based Anticancer Prodrugs as Potent Ferroptosis Inducers

Qianer Zhang,^{[a]#} Xuejing Fan,^{[a]#} Huimei Qian,^[a] Shanshan Xiao,^[a] Qin Song,^[a] Yicheng Wang,^[a] Jing Wang,^[a] Shuang Yang,^[a] and Yong Wang,^{*[a]}

[a] Key Laboratory of Marine Drugs, Chinese Ministry of Education; School of Medicine and Pharmacy, Ocean University of China, Qingdao 266003, Shandong, P.
R. China; Laboratory for Marine Drugs and Bioproducts, Qingdao Marine Science and Technology Center, Qingdao, 266237
E-mail: wangyong8866@ouc.edu.cn

[#] These two authors contributed equally

Table S1. Antitumor activities of representative NAAF prodrugs against HT1080
cells
Fig. S1. Expanded regions of the spectra about I-1 and II-1 with the generation of
boronic acid in the absence of H_2O_2 2
Fig. S2. Expanded regions of the spectra about I-1 hydrolysis in the presence of H_2O_2 .
Fig. S3. Expanded regions of the spectra about II-1 hydrolysis in the presence of
H ₂ O ₂ 5
Fig. S4. Expanded regions of the spectra about II-1-B hydrolysis in the presence of
H ₂ O ₂
Fig. S5. Activation of II-1 in HT1080 cell verified by LC-MS/MS
¹ H, ¹³ C NMR and HRMS spectra of compounds9
HPLC analysis for compounds



Table S1. Antitumor activities of representative NAAF prodrugs against HT1080 cells.

^{*a*} All values are expressed as the mean \pm SD from three independent experiments. ^{*b*} HT1080 and incubated with compounds alone. ^{*c*} HT1080 incubated with compounds and Fer-1 (1.5 μ M). ^{*d*} + Fer-1 IC₅₀/- Fer-1 IC₅₀.



Fig. S1. Expanded regions of the spectra of I-1 (top) and II-1 (bottom) with the generation of boronic acid in the absence of H_2O_2 .





Fig. S2. Expanded regions of the spectra of I-1 hydrolysis in the presence of H_2O_2 .



Fig. S3. Expanded regions of the spectra of II-1 hydrolysis in the presence of H_2O_2 .



Fig. S4. Expanded regions of the spectra of II-1-b hydrolysis in the presence of H_2O_2 .





Fig. S5. Activation of II-1 in HT1080 cell verified by LC-MS/MS.

¹H, ¹³C NMR and HRMS spectra of compounds

I-1 ¹H-NMR (Chloroform-*d*)



















I-1 HRMS

20230821-QR-9_230822093119 #53 RT: 0.58 AV: 1 NL: 1.42E7 T: FTMS + p ESI Full ms [180.00-2000.00]



I-2 HRMS



I-3 HRMS

20230821-QR-21_230822104709 #60 RT: 0.53 AV: 1 NL: 2.77E6 T: FTMS + p ESI Full ms [180.00-2000.00]



II-1 HRMS





II-1-B LCMS



II-2 HRMS

20230821-QR-10_230822093119 #47-48 RT: 0.39-0.40 AV: 2 NL: 1.05E6 T: FTMS + p ESI Full ms [180.00-2000.00]



III-1 HRMS





III-2 HRMS

20230821-QR-13 230822093119 #54 RT: 0.55 AV: 1 SB: 16 0.27-0.39 NL: 1.09E7 T: FTMS + p ESIFulims (180.00-2000.00]



HPLC analysis for compounds Compound I-1

Result

	Ret. Time	Area	Rel. Area (%)
1	2.576	176761	2.89
2	13.185	5900938	96.53
3	13.818	35208	0.58



Compound I-2

Result				
	Ret. Time	Area	Rel. Area (%)	
1	2.292	33829	1.34	
2	2.690	12662	0.50	
3	13.208	2462095	97.60	
4	13.765	7781	0.31	
5	14.117	6287	0.25	



Compound I-3

Result			
	Ret. Time	Area	Rel. Area (%)
1	14.066	4160493	97.59
2	14.672	32932	0.77
3	14.991	20228	0.47
4	15.286	49596	1.16



Compound II-1

Result

	Ret. Time	Area	Rel. Area (%)
1	11.076	14990	2.65
2	11.810	550991	97.35



Compound II-2

Resul	t
-------	---

	Ret. Time	Area	Rel. Area (%)
1	10.880	10367	0.88
2	13.819	1170541	99.12



Compound III-1

Resu	ılt

	Ret. Time	Area	Rel. Area (%)
1	10.113	3852405	95.65
2	10.399	41980	1.04
3	10.775	100230	2.49
4	11.660	33113	0.82



Compound III-2

Resul	t
-------	---

	Ret. Time	Area	Rel. Area (%)
1	13.981	3112944	98.36
2	14.482	51911	1.64

