

Supporting information for

Vortex Fluidic-Mediated Transesterification Enhancement of Mongongo Fatty Acid Ethyl Ester Production for Haircare Applications

Xuejiao Cao^a, Xuan Luo^a, Safira M. Barros^a, Wenjin Xing^a, Melanie MacGregor^a, Iliana Delcheva^a, Jonathan Campbell^a, Shan He^{a,b}, Bradley P. Kirk^c, Youhong Tang^a, Colin L. Raston^{a*}

^a Flinders Institute for Nanoscale Science and Technology, College of Science and Engineering, Flinders University, Adelaide, 5042, Australia

^b School of Food and Pharmacy, Zhejiang Ocean University, Zhejiang, 316022, China

^c Institute for Photonic and Advanced Sensing, School of Physics, Chemistry and Earth Science, University of Adelaide, South Australia 5005, Australia




Figure S1. Comparison of Mongongo FAEE processed via VFD and conventional method. a). VFD product after being washed with water. b). Traditional product of method after being washed with water.




Figure S2. Comparison of DSC of hair treated with Mongongo FAEE and Mongongo oil.




Figure S3. SEM-EDX of hair treated with Mongongo FAEE: C, N, O, and S elemental distribution on hair surface after the treatment.



hair.mp4

Vedio 1. Hair after 8 hours of sunlight treatment. From left to right, hair without oil/FAEE, hair with Mongongo oil, hair with Mongongo FAEE.