Supporting Information

Nitrogen Doped Carbon Quantum Dots: A Multifaceted Carbon Nanomaterial that interferes in an amyloid-forming trajectory

Sherin M. ElMorsy,[1] Denisse A Gutierrez,[2], Salvador Valdez,[3] Jyotish Kumar,[3] Renato J Aguilera,[2] Mohamed Noufal,[4] Sampath Chinnam,[5] Hemen Sarma,[6] and Mahesh Narayan[3,]*

[1]The Environmental Science & Engineering Program, The University of Texas at El Paso, El Paso, TX 79968, USA

[2]Cellular Characterization and Biorepository Core Facility, Border Biomedical Research Center, Department of Biological Sciences, College of Science, The University of Texas at El Paso, 500 West University Avenue, El Paso, TX 79968-0519, USA.

[3]Department of Chemistry and Biochemistry, The University of Texas at El Paso, 500 West University Avenue, El Paso, Texas79968, United States.

[4]Department of Chemical Engineering, Hampton University, Hampton, VA 23668, USA.

[5]Department of Chemistry, M.S. Ramaiah Institute of Technology (Autonomous Institution, Affiliated to Visvesvaraya Technological University, Belgaum), Bengaluru, Karnataka 560054, India

[6]Bioremediation Technology Research Group, Department of Botany, Bodoland University, Rangalikhata, Deborgaon, 783370, Kokrajhar (BTR), Assam, India

**Corresponding Author. E-mail: mnarayan@utep.edu. Mailing address: Dept. of Chemistry and Biochemistry, UTEP, 500 W. University Ave., El Paso, TX 79968. Phone: 915-747-6614. Fax: 915-747-5748.*

ORCID: Mahesh Narayan: 0000-0002-2194-5228

Table S1: Antioxidant activity comparison between N-CQDs and recently reported CQDs.

Carbon nanomaterials	DPPH Assay (Scavenging Efficiency)	Concentration	Reference
Graphene Quantum Dots	79%	20 μg/mL	[1]
Graphene oxide	42%	20 μg/mL	[1]
Se-CQDs	40%	200 μg/mL	[2]
Selenium and Nitrogen Co-Doped Carbon Quantum Dots	55%	0.4 mg/ml	[3]
Carbon dots	12.5 %	20 μg/ mL[1]	[4]
Nitrogen Doped Carbon Quantum Dots	41%	100 μg/ mL[1]	

References

(1) Hu, X.; Ma, X.-Y.; Tian, J.; Huang, Z. Rapid and facile synthesis of graphene quantum dots with high antioxidant activity. *Inorganic Chemistry Communications* **2020**, *122*, 108288. DOI: https://doi.org/10.1016/j.inoche.2020.108288.

(2) Huang, G.; Lin, Y.; Zhang, L.; Yan, Z.; Wang, Y.; Liu, Y. Synthesis of Sulfur-Selenium Doped Carbon Quantum Dots for Biological Imaging and Scavenging Reactive Oxygen Species. *Scientific Reports* **2019**, *9* (1), 19651. DOI: 10.1038/s41598-019-55996-w.

(3) Xu, Q.; Liu, X.; Jiang, Y.; Wang, P. A highly sensitive and selective probe for the colorimetric detection of Mn (II) based on the antioxidative selenium and nitrogen co-doped carbon quantum dots and ABTS•+. *Frontiers in Chemistry* **2021**, *9*, 658105.

(4) Innocenzi, P.; Stagi, L. Carbon dots as oxidant-antioxidant nanomaterials, understanding the structure-properties relationship. A critical review. *Nano Today* **2023**, *50*, 101837. DOI: https://doi.org/10.1016/j.nantod.2023.101837.